

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 391 516 A1**

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 158(3) EPC

(43) Date of publication:
25.02.2004 Bulletin 2004/09

(51) Int Cl.7: **C12N 15/57**

(21) Application number: **02722783.4**

(86) International application number:
PCT/JP2002/004141

(22) Date of filing: **25.04.2002**

(87) International publication number:
WO 2002/088366 (07.11.2002 Gazette 2002/45)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

- **MIMURA, Noriko, c/o Kikuchi Research Center
Kumamoto 869-1298 (JP)**
- **MAEDA, Hiroaki, c/o Kikuchi Research Center
Kumamoto 869-1298 (JP)**
- **NOZAKI, Chikateru, c/o Kikuchi Research Center
Kumamoto 869-1298 (JP)**
- **HAMAMOTO, Takayoshi
Kumamoto-shi, Kumamoto 860-8568 (JP)**
- **NAKAGAKI, Tomohiro
Kumamoto-shi, Kumamoto 860-8568 (JP)**

(30) Priority: **25.04.2001 JP 2001128342**
27.07.2001 JP 2001227510
28.09.2001 JP 2001302977
25.01.2002 JP 2002017596

(71) Applicant: **Juridical Foundation,
The Chemo-Sero-Therapeutic Research Institute
Kumamoto-shi, Kumamoto 860-8568 (JP)**

(74) Representative: **HOFFMANN - EITLE
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)**

(72) Inventors:
• **SOEJIMA, Kenji, c/o Kikuchi Research Center
Kumamoto 869-1298 (JP)**

(54) **VON WILLEBRAND FACTOR (VWF)-CLEAVING ENZYME**

(57) This invention is intended to isolate and identify a vWF-specific cleaving protease.

The vWF-specific cleaving protease cleaves a bond between residues Tyr 842 and Met 843 of vWF and comprises a polypeptide chain having Leu-Leu-Val-Ala-Val as a partial sequence, and more preferably comprises a polypeptide chain having the partial N-terminal amino acid sequence of a mature protein, Ala-Ala-Gly-Gly-Ile-

Leu-His-Leu-Glu-Leu-Leu-Val-Ala-Val, and having a molecular weight of 105 to 160 kDa in SDS-PAGE under reducing or non-reducing conditions. Isolation and identification of this vWF-specific cleaving protease have led to the possibility of replacement therapy for patients having diseases resulting from a deficiency of the protease, such as thrombotic thrombocytopenic purpura.

EP 1 391 516 A1

Description

Technical Field

5 [0001] The present invention relates to a plasma protein related to the field of medical drugs. More particularly, the present invention relates to a protease that specifically cleaves von Willebrand factor (it may be hereafter referred to as "vWF"), which is associated with blood coagulation. The vWF-cleaving protease of the present invention enables replacement therapy for patients with diseases resulting from defects or decreases in this protease, such as thrombotic thrombocytopenic purpura (it may be hereafter referred to as "TTP"). In addition, the use thereof as a novel antiplatelet

10 thrombotic agent is expected.

Background Art

15 [0002] vWF is produced in vascular endothelial cells or megakaryocytes, and is a blood coagulation factor in which a single subunit comprising 2,050 amino acid residues (monomers of about 250 kDa) are bound by an S-S bond to form a multimer structure (with a molecular weight of 500 to 20,000 kDa). The level thereof in the blood is about 10 µg/ml, and a high-molecular-weight factor generally has higher specific activity.

[0003] vWF has two major functions as a hemostatic factor. One of the functions is as a carrier protein wherein vWF binds to the blood coagulation factor VIII to stabilize it. Another function is to form platelet plug by adhering and agglomerating platelets on the vascular endothelial subcellular tissue of a damaged vascular wall.

20 [0004] Thrombotic thrombocytopenic purpura is a disease that causes platelet plug formation in somatic arterioles and blood capillaries throughout the whole body. In spite of recent advances in medical technology, the morbidity associated with this disease approximately tripled from 1971 to 1991. Pathologically, TTP is considered to result from vascular endothelial cytotoxicity or vascular platelet aggregation. Immunohistologically, a large amount of vWFs are recognized in the resulting platelet plugs, and vWF is considered to play a major role in causing them. A normal or high-molecular-weight vWF multimer structure is dominant in a TTP patient, and an unusually large vWF multimer (ULvWFM) or large vWF multimer (LvWFM) is deduced to play a major role in accelerating platelet aggregation or microthrombus formation under high shearing stress. In contrast, vWF was known to degrade at a position between residues Tyr 842 and Met 843 by the action of vWF-cleaving protease in the circulating blood of a healthy person under

25 high shearing stress. Accordingly, TTP is considered to occur in the following manner. The protease activity in the plasma is lowered for some reason, and ULvWFM to LvWFM are increased to accelerate platelet aggregation. This forms platelet plugs in blood vessels.

30 [0005] Recently, Furlan et al. (Blood, vol. 87, 4223-4234: 1996, JP Patent Publication (Kohyo) No. 2000-508918) and Tsai et al. (Blood, vol. 87, 4235-4244: 1996) developed a method for assaying vWF-specific cleaving protease. In their report, this protease activity was actually lowered in TTP. The aforementioned authors reported that this enzyme was metalloprotease in the plasma and partially purified. However, they have not yet succeeded in the amino acid sequencing which would specify the protease. There have been no further developments since then.

35

Disclosure of the Invention

40 [0006] Up to the present, plasmapheresis therapy has been performed for treating patients who congenitally lack vWF-specific cleaving protease and patients who had acquired positive antibodies against this protease. Establishment of replacement therapy using purified products or a pure substance such as a recombinant gene product of the aforementioned protease is desired. Familial TTP patients congenitally lack vWF-specific cleaving protease, and non-familial

45 TTP is caused by posteriori production of autoantibodies against the aforementioned protease. Accordingly, replacement therapy for this protease is preferable for familial TTP patients (plasma administration is actually performed), and removal of autoantibodies by plasmapheresis and substitution of this protease are necessary for non-familial TTP. Further, the use of this protease as a novel antiplatelet thrombotic agent can also be expected.

50 [0007] As mentioned above, however, Furlan et al. (Blood, vol. 87, 4223-4234: 1996, JP Patent Publication (Kohyo) No. 2000-508918) and Tsai et al. (Blood, vol. 87, 4235-4244: 1996) have suggested that the vWF-cleaving protease was metalloprotease in the plasma. It was reported to be partially purified, and concentrated 1,000- to 10,000-fold from the plasma in terms of its specific activity. Even under these conditions, there has been no advancement in the analysis of the properties of this protease, such as the amino acid sequence of its protein, over the period of roughly 5 years that has passed since then. No specific biological information has yet been obtained regarding this protease. As reported by Furlan et al., the protein of interest is supposed to be gigantic, and there may be various problems associated therewith. For example, diversified forms of this protease, such as various interacting molecules or cofactors, are expected. Based on the complexity of purification processes, deteriorated capacity of separation by nonspecific inter-

55 action during the purification step, and other factors, it is deduced to be very difficult to isolate and identify the protease

from a plasma fraction by the purification process according to Furlan et al.

[0008] Under the above circumstances, the present inventors have conducted concentrated studies in order to isolate and identify the vWF-cleaving protease. As a result, they have succeeded in isolating and purifying the vWF-cleaving protease of interest, which had not yet been reported. Thus, they have succeeded in identifying an amino acid sequence of the mature protein and a gene encoding this amino acid sequence.

[0009] The vWF-cleaving protease of the present invention can cleave a bond between residues Tyr 842 and Met 843 of vWF. According to one embodiment, this protease has a molecular weight of 105 to 160 kDa or 160 to 250 kDa in SDS-PAGE under reducing or non-reducing conditions. It is comprised of a polypeptide chain having Leu-Leu-Val-Ala-Val as a partial sequence. More preferably, it is comprised of a polypeptide chain having the partial N-terminal amino acid sequence of a mature protein, i.e., Ala-Ala-Gly-Gly-De-Leu-His-Leu-Glu-Leu-Leu-Val-Ala-Val. It is a novel substance characterized by the following properties.

1) vWF-cleaving activity

[0010] According to the N-terminal sequence analysis of the cleavage fragment, the protease of the present invention cleaves a peptide bond between residues Tyr 842 and Met 843.

2) Fractionation by gel filtration

[0011] When fractionation is performed by gel filtration chromatography using FI paste as a starting material, most activities are collected in a fraction with a molecular weight of 150 to 300 kDa. According to one embodiment of the present invention, an actually obtained active substance is found to have a molecular weight of about 105 to 160 kDa in electrophoresis. Accordingly, the protease of the present invention is a substance that is likely to form a dimer or the like or to bind to another molecule or a substance that can be easily degraded or can have a heterogeneous sugar chain added.

3) Ammonium sulfate precipitation

[0012] For example, when FI paste is used as a starting material, a large portion of this protease is recovered as a precipitation fraction from a roughly purified fraction with the use of 33% saturated ammonium sulfate.

4) SDS-PAGE

[0013] For example, the protease of the present invention derived from FI paste prepared from pooled human plasma or cryoprecipitate mainly has a molecular size of about 105 to 160 kDa determined by a molecular weight marker in SDS-PAGE. Based on the nucleic acid sequence as shown in SEQ ID NO: 15, when an amino acid sequence represented by a frame between an atg initiation codon at position 445 and a tga termination codon at position 4726 is expressed by gene recombination, there are some variations in molecular sizes depending on a host. However, a molecular size of about 160 to 250 kDa determined by a molecular weight marker is exhibited. This size is observed in the plasma of healthy humans and in that of some TTP patients. Several molecular species of this protease are present in human plasma, caused by the presence of alternative splicing products (SEQ ID NOs: 16 to 21) recognized at the time of gene cloning, differences in post-translational modification such as sugar chain addition, or degradation during purification. Further, this protease could be partially recovered in an active state after SDS-PAGE under non-reducing conditions.

5) Analysis of amino acid sequence

[0014] The amino acid sequence of the isolated polypeptide fragment was analyzed. This presented an example of a polypeptide chain having a sequence Leu-Leu-Val-Ala-Val as a partial amino acid sequence and a sequence Ala-Ala-Gly-Gly-Ile-Leu-His-Leu-Glu-Leu-Leu-Val-Ala-Val as a N-terminal amino acid sequence of a mature protein. Further, with current bioinformatics (BIOINFORMATICS: A Practical Guide to the Analysis of Genes and Proteins, edited by Andreas D. Baxeavanis and B. F. Francis Ouellette), a nucleic acid sequence encoding the amino acid sequence was highly accurately identified by searching a database based on the aforementioned partial sequence. More specifically, the genome database was searched by the tblastn program. This identified a chromosome clone (AL158826) that is deduced to encode the protease of the present invention. Further, clones (AI346761 and AJ011374) that are deduced to be a part of the protease of interest and a part of the polypeptide to be encoded by the aforementioned genome were identified through collation with the Expressed Sequence Tag (EST) database. Based thereon, the amino acid sequence as shown in SEQ ID NO: 3 or 7 was identified as an active vWF-cleaving protease site.

[0015] GCT GCA GGC GGC ATC CTA CAC CTG GAG CTG CTG GTG GCC GTG, a sequence deduced from the genome, and more preferably CTG CTG GTG GCC GTG, a portion thereof, the transcriptome of which was confirmed by EST, was obtained. The obtained nucleotide sequence was analyzed, and motif analysis was carried out based on the deduced sequence. As a result, it was found to have a metalloprotease domain as a candidate for the protease of the present invention. Based on the above findings, it became possible to disclose a sequence of a polypeptide chain as a more specific example of the protease. Also, activities of proteases are generally known to vary depending on, for example, substitution, deletion, insertion, or introduction of point mutation into a portion of the amino acid sequence (Blood coagulation factor VII mutants, Soejima et al., JP Patent Publication (Kokai) No. 2001-61479 A). Similarly, the protease of the present invention, can be modified by, for example, deletion, substitution, or addition of one or several amino acids, to prepare optimized proteases.

[0016] The protease proteins were further mass-produced, and 29 amino acid sequences from the N-terminus were determined. These amino acid sequences are shown in SEQ ID NO: 8. This result is substantially the same as the sequence as shown in SEQ ID NO: 3 or 7 deduced by bioinformatics. Only one difference is that the amino acid 27th in SEQ ID NO: 3 or 7 was Glu while it was Arg according to the present analysis of the N-terminal sequence. This was considered to be a gene polymorphism. Thus, this protease was confirmed to be comprised of a polypeptide chain having the amino acid sequence as shown in SEQ ID NO: 3 or 7 at its N-terminus as a mature unit. A gene fragment encoding this protease was then cloned in the following manner.

[0017] Based on the nucleic acid sequence as shown in SEQ ID NO: 7, a sense primer (SEQ ID NO: 9) and an antisense primer (SEQ ID NO: 10) were prepared based on the nucleic acid sequence underlined in Fig. 9, and a gene sandwiched between these primers was amplified. This fragment was cloned, and the nucleotide sequence was then confirmed. This fragment was used as a probe for Northern blotting to analyze the site at which the protease gene was expressed. As a result, this protease gene was found to be expressed mainly in the liver. Accordingly, the human liver cDNA library was purchased, and a gene encoding this protease was identified using a rapid amplification of cDNA ends (RACE) technique. Based on these results, in the case of the largest sequence of approximately 5 kb of mRNA (cDNA) reaching the poly(A) addition site as shown in SEQ ID NO: 15 was identified.

[0018] Based on the amino acid sequence deduced from this gene sequence, this protease was deduced to have a preprosequence, and to belong to the disintegrin and metalloprotease (ADAM) family having a disintegrin-like domain, a metalloprotease domain, and the like, and particularly to the ADAM-TS family having a thrombospondin Type-1 (TSP-1) domain. Finally, including those having insertion or deletion in a part of the nucleic acid sequence, isoforms as shown in SEQ ID NOs: 16 to 21 having sequences as shown in SEQ ID NOs: 3 and 7 at the N-terminuses after the mature preprosequence has been cleaved were identified. Thus, the protease of the present invention should cleave vWF between residues Tyr 842 and Met 843 and should have the Leu-Leu-Val-Ala-Val sequence as a partial amino acid sequence.

[0019] The vWF-cleaving protease of the present invention can be generally prepared by the following process.

[0020] According to the present invention, a process for assaying the protease activity is characterized by the possibility of evaluating activity within a short period of time. According to the report by Furlan et al. (Blood, vol. 87, 4223-4234: 1996, JP Patent Publication (Kohyo) No. 2000-508918 A), activity is assayed by analyzing vWF-cleaving patterns by Western blotting using the anti-vWF antibody, and thus, it takes time to transfer the protease to a filter. More specifically, this process requires approximately at least 45 hours in total, i.e., 24 hours for the enzymatic reaction with a substrate vWF, 17 hours for electrophoresis, and 3 hours to transfer the protease to a filter, followed by detection using the anti-vWF antibody. In contrast, the present inventors completed activity assay in 18 hours in total, i.e., 16 hours for the enzymatic reaction with a substrate vWF, and 2 hours for electrophoresis and detection. This indicates that the time required for the assay can be reduced to one third or less of that required for the conventional assay. This can also shorten the time required for the purification process, and in turn can lower the degree of the protease to be inactivated. Accordingly, purification efficiency is improved compared with that attained by the method of Furlan et al., and as a result, the degree of purification is also enhanced.

[0021] Further, the starting material was examined using the aforementioned assay system. As a result, it was found that the protease activity was more concentrated in FI paste than in the cryoprecipitate that had been reported by Furlan et al. in the past. FI paste was used as a starting material, and the aforementioned rapid activity assay systems were combined. This enabled isolation and identification of the protease of interest. In a specific embodiment, a purification process combining gel filtration chromatography with ion exchange chromatography is employed, and the aforementioned activity assay system is also combined.

[0022] More specifically, FI paste is solubilized with a buffer, and the resultant is fractionated by gel filtration chromatography. The protease activity is fractionated at the elution region with a molecular weight of 150 to 300 kDa deduced from the size marker of gel filtration. Thereafter, the resultant is precipitated and concentrated using 33% saturated ammonium sulfate. This procedure is repeated three times in total. The active fraction obtained in the third gel filtration is pooled, and the resultant is subjected to dialysis at 4°C overnight with a buffer comprising 50 mM NaCl added to 50 mM Tris-HCl (pH 7.1). Thereafter, the dialysis product is subjected to anion exchange chromatography

(DEAE) and eluted stepwise with 0.25 M NaCl. The present inventors have conducted concentrated studies in order to find a process for isolating and identifying the protease of the present invention. As a result, they found that, surprisingly, the protease was recoverable as an active band after non-reducing SDS-PAGE. In order to achieve further mass production, the purified and concentrated fraction was applied to the Biophoresis utilizing the principle of SDS-PAGE. Thus, a fraction having vWF-cleaving activity was isolated from the electrophoresed fraction. According to the approximate calculation of the specific activity up to this phase, purification of about 30,000- to 100,000-fold was achieved. This procedure was efficiently and rapidly repeated several times, and thus, about 0.5 pmole of sample that is the current limit of the analysis of amino acid sequence was obtained. Thus, analysis of amino acid sequence became feasible. More specifically, a final step of separation and purification (Biophoresis) based on the principle of SDS-PAGE is important, and it is based on the findings as a result of concentrated studies, which had led to the completion of the present invention.

[0023] According to the report by Furlan et al., specific activity was improved by as much as about 10,000 times, although the protease was not substantially isolated or identified. This could be because of deactivation during purification or the difficulty of isolating and identifying molecules, which were gigantic proteins capable of interacting with various other proteins such as the protease of the present invention by a separation method utilizing various types of liquid chromatography. Further, the protease content in the plasma was deduced to be very small, and thus, it was necessary to await the establishment of the process according to the present invention. Furthermore, the use of this process enables the purification of recombinant genes.

[0024] Based on the findings of the present invention, peptides or proteins prepared from the obtained sequences are determined to be antigens. With the use thereof, a monoclonal antibody, a polyclonal antibody, or a humanized antibody thereof can be prepared by general immunization techniques (Current Protocols in Molecular Biology, Antibody Engineering: A PRACTICAL APPROACH, edited by J. McCafferty et al. or ANTIBODY ENGINEERING second edition, edited by Carl A. K. BORREBAECK). Alternatively, an antibody that binds to the aforementioned protein can be prepared by antibody-producing techniques utilizing phage display (Phage Display of Peptides and Proteins: A Laboratory Manual, edited by Brian K. Kay et al., Antibody Engineering: A PRACTICAL APPROACH, edited by J. McCafferty et al. or ANTIBODY ENGINEERING second edition, edited by Carl A. K. BORREBAECK). Alternatively, based on these techniques, a neutralizing antibody acting against the protease activity or a simple binding antibody can be isolated from a specimen from a TTP patient who has an autoantibody positive against this protease. These antibodies can be applied to diagnosis and therapy of diseases such as TTP.

[0025] Based on the obtained genome or EST sequence, cDNA or a genomic gene encoding the protease of the present invention can be cloned by a common technique (Molecular Cloning, 2nd edition). Further, bioinformatics techniques (BIOINFORMATICS: A Practical Guide to the Analysis of Genes and Proteins, edited by Andreas D. Baxevanis and B. F. Francis Ouellette) enable cloning of the proteins of other animal species that are homologous thereto, and the resultant gene is fractured by a common technique (for example, Gene Targeting: A Practical Approach, First Edition, edited by A. L. Joyner, Teratocarcinomas and embryonic stem cell a practical approach) to produce TTP-like animal models. In particular, the identification of the gene sequence encoding the protein derived from a mouse enables the production of a knockout mouse having this gene. Thus, a disease mouse model of congenital TTP or the like can be prepared.

[0026] In accordance with a common technique (for example, J. Sambrook et al., Molecular Cloning, 2nd edition, or CURRENT PROTOCOLS IN MOLECULAR BIOLOGY), these genes are incorporated into a suitable expression vector, the resultant is transformed into a suitable host cell, and the gene recombinant product of the protease can be thus prepared. In this case, the gene to be incorporated is not necessarily the one that encoded the entire region of the protein. It also includes a partial expression of the protein as defined by a domain depending on its usage.

[0027] For example, the polynucleotide according to the present invention is introduced into a host cell using a conventional technique such as transduction, transfection, or transformation. The polynucleotide is introduced solely or together with another polynucleotide. Another polynucleotide is introduced independently, simultaneously, or in combination with the polynucleotide of the present invention.

[0028] For example, the polynucleotide of the present invention is transfected in a host cell, such as a mammalian animal cell, by a standard technique for simultaneous transfection and selection using another polynucleotide encoding a selection marker. In this case, the polynucleotide would be generally stably incorporated in the genome of the host cell.

[0029] Alternatively, the polynucleotide may be bound to a vector comprising a selection marker for multiplication in a host. A vector construct is introduced to a host cell by the aforementioned technique. In general, a plasmid vector is introduced as DNA of a precipitate, such as a calcium phosphate precipitate, or a complex with a charged lipid. Electroporation is also employed for introducing the polynucleotide into a host. When the vector is a virus, this virus is packaged *in vitro* or introduced into a packaging cell, thereby introducing the packaged virus into a cell.

[0030] Extensive techniques that are suitable for producing a polynucleotide and introducing the resulting polynucleotide to a cell in accordance with this embodiment of the present invention are known and common in the art. Such techniques are described in Sambrook et al. (aforementioned), and this document explains a variety of standard 'ex-

perimental manuals describing the aforementioned techniques in detail. In respect of this embodiment of the present invention, the vector is, for example, a plasmid vector, a single- or double-stranded phage vector, or a single- or double-stranded RNA or DNA viral vector. Such a vector is introduced into a cell as a polynucleotide, and preferably as DNA by a common technique for the introduction of DNA or RNA into a cell. When the vector is a phage or virus, the vector is preferably introduced to the cell as a packaged or sealed virus by a known technique for infection and transduction. A viral vector may be of a replication-competent or defective type.

[0031] A preferable vector is a vector which expresses the polynucleotide or polypeptide of the present invention in points. In general, such a vector comprises a cis-action control region that is effective for the expression in a host operably bound to the polynucleotide to be expressed. When a suitable trans-action factor (for example, a group of proteases involved with the post-translational processing such as signal peptidase or Furin) is introduced in a host cell, it is supplied by a host, a complementary vector, or the vector itself.

[0032] In a preferable embodiment, a vector provides specific expression. Such specific expression is an inducible one or realized only in a certain type of cell. Alternatively, it is an inducible and cell-specific expression. A particularly preferable inducible vector can induce expression by an easily operable environmental factor such as temperature or a nutritional additive. Various vectors suitable for this embodiment including a construction for the use in prokaryotic and eukaryotic cell hosts and an inducible expression vector are known, and persons skilled in the art can commonly use them.

[0033] A genetically engineered host cell can be cultured in general nutrient medium, and it is modified to be particularly suitable for activation of promoter, selection of transformant, or amplification of a gene. In general, it would be obvious to persons skilled in the art that conventional culture conditions such as temperature or pH level for host cells selected for the expression are suitable for the expression of the polypeptide of the invention.

[0034] A wide variety of expression vectors can be used for expressing the polypeptide of the present invention. Examples of these vectors include chromosome, episome, and virus-derived vectors. These vectors are derived from bacterial plasmid, bacteriophage, yeast episome, yeast chromosome element, or viruses such as baculovirus, papovavirus such as simian virus 40 (SV40), vaccinia virus, adenovirus, fowlpox virus, pseudorabies virus, or retrovirus. A vector derived from a combination of the aforementioned, for example, a vector derived from plasmid and bacteriophage gene element, more specifically, a cosmid or phagemid, may also be used. They are used for the expression in accordance with this embodiment of the present invention. In general, since polypeptides were expressed in hosts, any vector that is suitable for maintaining, multiplying, or expressing a polynucleotide can be used for the expression according to the aforementioned embodiment. A suitable DNA sequence is inserted into a vector by various conventional techniques. In general, a DNA sequence for expression is bound to an expression vector by cleavage of a DNA sequence and an expression vector having 1 or more restriction endonucleases, and a restriction fragment is then bound together using T4 DNA ligase. Restriction and ligation techniques that can be used for the above purpose are known and common to persons skilled in the art. With regard thereto, Sambrook et al. (aforementioned) very precisely describe another suitable method for constructing an expression vector utilizing another technique known and common to persons skilled in the art.

[0035] A DNA sequence in the expression vector is operably bound to, for example, a suitable expression-regulating sequence including a promoter to orient the mRNA transcription. A few examples of known representative promoters are the phage lambda PL promoter, *E. coli* lac, trp, trc, and tac promoters, SV40 early and late promoters, and the retrovirus LTR promoter. Many promoters that are not described are suitable for the use according to the embodiment of the present invention, known, and more easily used as described in the examples of the present invention. In general, an expression construct comprises a ribosome binding site for translation in a transcription initiation or termination site or a transcribed domain. The coding region of the mature transcript that was expressed by the construct comprises the initiation AUG at the initiation and termination codons located substantially at the terminus of polypeptide to be translated. In addition, the construct comprises a regulator region that regulates and induces the expression. In general, such a region is activated through the regulation of the repressor binding site, transcription of an enhancer, or the like in accordance with various conventional methods.

[0036] Vectors for multiplication and expression include selection markers. Such markers are suitable for multiplication, or they comprise additional markers for the above-stated purpose. The expression vector preferably comprises one or more selection marker genes to provide phenotypic traits for the purpose of selecting the transformed host cell. A preferable marker includes dihydrofolate reductase- or neomycin-resistance with regard to eukaryotic cell culture. It has tetracycline- or ampicillin-resistance with regard to *E. coli* and other bacterial cultures. A suitable vector comprising a DNA sequence and a suitable promoter or regulatory sequence as described herein are introduced to a suitable host by various suitable known techniques for the expression of the polypeptide of interest.

[0037] Representative examples of suitable hosts include: bacterial cells such as *E. coli*, *Streptomyces*, and *Salmonella typhimurium*; fungal cells such as a yeast cell; insect cells such as drosophila S2 and Spodoptera Sf9 cells; and adhesive or floating animal or plant cells such as CHO, COS, Bowes melanoma cells, and SP2/0. Various hosts for expression constructs are known, and persons skilled in the art can easily select a host for expressing polypeptides

in accordance with this embodiment based on the disclosure of the present invention.

[0038] More specifically, the present invention includes a recombinant construct, such as an expression construct comprising one or more sequences as mentioned above. The construct is a vector, such as a plasmid or viral vector comprising the sequence of the present invention inserted therein. The sequence is inserted in a positive or negative direction. In a preferable specific example thereof, the construct further has a regulatory sequence comprising a promoter or the like that is operably bound to the sequence. Various suitable vectors and promoters are known to persons skilled in the art, and there are many commercially available vectors that are suitably used in the present invention.

[0039] Commercially available vectors are exemplified below. Vectors that are preferably used for bacteria are pQE70, pQE60, and pQE-9 (Qiagen); pBS vector, PhageScript vector, Bluescript vector, pNH8A, pNH16a, pNH18A, and pNH46A (Stratagene); and ptrc99a, pKK223-3, pKK233-3, pDR540, and pRIT5 (Pharmacia). Examples of preferable eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1, and pSG (Stratagene) and pSVK3, pBPV, pMSG, and pSVL (Pharmacia). These vectors are commercially available for persons skilled in the art to be used in accordance with the embodiment of the present invention, and they are merely a list of known vectors. For example, other plasmids or vectors suitable for introducing, maintaining, multiplying, or expressing the polynucleotide or polypeptide of the present invention can also be used in hosts in accordance with this embodiment of the present invention.

[0040] A promoter region can be selected from a gene of interest using a vector comprising, for example, a candidate promoter fragment, i.e., a reporter transcription unit lacking a promoter region such as a chloramphenicol acetyl transferase (CAT) transcription unit located downstream of restriction sites for introducing promoter-containing fragments. As known to the public, the introduction of the promoter-containing fragment into the vector at the restriction site located upstream of the cat gene generates CAT activity that can be detected by standard CAT assay. A vector that is suitable for this purpose is known and readily available. Examples of such vectors are pKK232-8 and pCM7. Accordingly, the promoter for expressing the polynucleotide of the present invention includes not only a readily available known promoter but also a promoter that can be readily obtained using a reporter gene in accordance with the aforementioned technique.

[0041] Among them, according to the present invention, examples of known bacterial promoters that are suitably used to express polynucleotides and polypeptides are *E. coli* lacI and lacZ promoters, T3 and T7 promoters, gpt promoter, lambda PR and PL promoters, and trp and trc promoters. Examples of suitable known eukaryotic promoters include the Cytomegalovirus (CMV) immediate promoter, the HSV thymidine kinase promoter, early and late SV40 promoters, a retrovirus LTR promoter such as the Rous sarcoma virus (RoSV) promoter, and a metallothionein promoter such as the metallothionein-I promoter.

[0042] Selection of a vector and a promoter suitable for expression in a host cell is a known technique. Techniques necessary for the construction of expression vectors, introduction of a vector in a host cell, and expression in a host are common in the art. The present invention also relates to a host cell having the aforementioned construct. A host cell can be a higher eukaryotic cell such as a mammalian animal cell, a lower eukaryotic cell such as a yeast cell, or a prokaryotic cell such as a bacterial cell.

[0043] The construct can be introduced in a host cell by calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. These methods are described in a variety of standard laboratory manuals, such as a book by Sambrook et al.

[0044] The construct in a host cell can be used by a conventional method, and it produces a gene product encoded by a recombinant sequence. Alternatively, a partial polypeptide of the present invention can be synthesized using a general peptide synthesizer. A mature protein can be expressed under the control of a suitable promoter in a mammalian animal, yeast, bacterial, or other cell. Also, such a protein can be produced in a cell-free translation system with the use of RNA derived from the DNA construct of the present invention. Suitable cloning and expression vectors for prokaryotic and eukaryotic hosts are described in Sambrook et al (aforementioned).

[0045] In general, a recombinant expression vector comprises: a replication origin; a promoter derived from a highly expressed gene to orient the transcription of a downstream structural sequence; and a selection marker for bringing the cell into contact with a vector and isolating the vector-containing cell. A suitable promoter can be induced from a gene encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, and heat shock protein. A selection marker includes *E. coli* ampicillin-resistant gene and *S. cerevisiae* trp1 gene.

[0046] Transcription of DNA encoding the polypeptide of the present invention using a higher eukaryotic cell may be enhanced by inserting an enhancer sequence in a vector. The enhancer is generally a cis-acting element for DNA for enhancing the promoter transcription activity in the predetermined host cell. Examples of an enhancer include the SV40 enhancer, the Cytomegalovirus early promoter/enhancer, the polyoma enhancer behind the replication origin, the β -actin enhancer, and the adenovirus enhancer.

[0047] The polynucleotide of the present invention encoding a heterologous structural sequence of the polypeptide of the present invention is generally inserted in a vector by standard techniques in such a manner that it is operably bound to the expression promoter. The transcription initiation site of the polypeptide is suitably located at the 5' site of the ribosome binding site. The ribosome binding site is 5' relative to AUG that initiates the translation of a polypeptide to be expressed. In general, an initiation codon starts from AUG and another open reading frame located between the

ribosome binding site and initiation AUG is not present. The termination codon is generally present at the terminus of the polypeptide, and the adenylation signal and the terminator are suitably located at the 3' end of the transcription region.

[0048] Regarding the secretion of the translated protein in the ER lumen, in the cytoplasm, or to the extracellular environment, a suitable secretion signal is incorporated in the expressed polypeptide. The signal may be endogenous or heterologous to the polypeptide.

[0049] Further, a prosequence subsequent to the signal sequence may be endogenous or heterologous (e.g., a preprosequence of another metalloprotease).

[0050] The polypeptide is expressed in a modified form such as a fusion protein, and it includes not only a secretion signal but also an additional heterologous functional region. Accordingly, an additional amino acid, especially a charged amino acid region, or the like, is added to the polypeptide to improve stability and storage stability in the host cell during purification or subsequent operation and storage. Alternatively, a given region may be added to the polypeptide to accelerate the purification. This type of region may be removed before the final preparation of polypeptides. Induction of secretion or excretion, stability improvement, or facilitation of purification with the addition of a peptide portion to the polypeptide is a technique common and known in the art.

[0051] Examples of prokaryotic hosts that are suitable for multiplying, maintaining, or expressing the polynucleotide or polypeptide of the present invention include *E. coli*, *Bacillus subtilis*, and *Salmonella typhimurium*. Various types of *Pseudomonas*, *Streptomyces*, and *Staphylococcus* are suitable hosts in this respect. Furthermore, various other types of hosts known to persons skilled in the art can be also used. Representative examples of expression vectors that are useful for bacterial applications include, but are not limited to, the replication origin of bacteria derived from commercially available plasmid including a selectable marker and a gene element of a known cloning vector pBR322 (ATCC 37017). Examples of such commercially available vectors include pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, Wisconsin, USA). These pBR322 (main chain) sections are combined with a suitable promoter and structural sequences to be expressed.

[0052] Host cells are suitably transformed and multiplied to the optimal cell concentration. Thereafter, the selected promoter is induced by a suitable means (e.g., temperature shifting or chemical inducer), and cells are further cultured. Typically, cells are collected by centrifugation and fractured by a physical or chemical means. The resulting crude extract is further purified. Microbial cells used for the protein expression can be fractured by any convenient means selected from a freezing-thawing cycle, ultrasonication, mechanical fracture, and the use of a cytolytic agent. These methods are known to persons skilled in the art.

[0053] Various cell lines for mammalian animal cell culture can be also used for the expression. An example of a cell line for mammalian animal expression includes a monkey kidney fibroblast COS-6 cell described in Gluzman et al., Cell 23: 175 (1981). Examples of other cells that are capable of expressing compatible vectors include C127, 3T3, CHO, HeLa, human kidney 293, and BHK cells. Further, a floating myeloma cell line such as SP2/0 can be also used.

[0054] A mammalian animal expression vector comprises a replication origin, a suitable promoter and enhancer, a necessary ribosome binding site, a polyadenylation site, splice donor and acceptor sites, a transcription termination sequence, and a 5' flanking untranscribed sequence necessary for expression. DNA sequences derived from the SV40 splice site and the SV40 polyadenylation site are used for the non-transformed or transcribed gene element of interest. An example thereof is a CAG expression vector (H. Niwa et al., Gene, 108, 193-199 (1991)).

[0055] Based on the gene sequence of the above protease, a probe, primer, or antisense is designed by a common technique. The antisense technique can be used for controlling gene expression by the use of antisense DNA or RNA or the formation of a triple helix. This technique is described in, for example, Okano, J., Neurochem., 56: 560 (1991); OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION, CRC Press, Boca Raton, FL (1988). The triple helix formation is examined in, for example, Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). The method is based on the polynucleotide bond with complementary DNA or RNA. This enables the gene diagnosis or gene therapy.

[0056] For example, cells obtained from a patient are subjected to *ex vivo* genetic engineering using a polynucleotide such as polypeptide-encoding DNA or RNA. The resulting cells are then supplied to patients who should be treated with polypeptides. For example, cells can be subjected to *ex vivo* genetic engineering using a retrovirus plasmid vector comprising RNA encoding the polypeptide of the present invention. Such a technique is known in the art, and the use thereof in the present invention is obvious according to the description given herein. Similarly, cells are subjected to *in vitro* genetic engineering in accordance with a conventional process in respect of *in vivo* polypeptide expression. For example, the polynucleotide of the present invention is genetically engineered for expression in the replication-deficient retrovirus vector as mentioned above. Subsequently, the retrovirus expression construct is isolated, introduced to a packaging cell, and transduced using a retrovirus plasmid vector comprising RNA encoding the polypeptide of the present invention. Thus, the packaging cell produces infectious viral particles having a control gene. These producer cells are subjected to *in vitro* genetic engineering and then administered to patients to allow polypeptides to be expressed *in vivo*. This administration method and other methods for administering polypeptides according to the present

invention would be clearly understood by persons skilled in the art based on the teaching of the present invention.

[0057] Examples of the aforementioned retrovirus, from which the retrovirus plasmid vector is derived, include, but are not limited to, Moloney murine leukemia virus, spleen necrosis virus, Rous sarcoma virus, Harvey sarcoma virus, avian leukosis virus, gibbon leukemia virus, human immunodeficiency virus, myeloproliferative sarcoma virus, and mammary tumor virus. This type of vector comprises one or more promoters to express polypeptides. Examples of suitable promoters that can be used include, but are not limited to, retrovirus LTR, SV40 promoter, CMV promoter described in Miller et al., *Biotechniques* 7: 980-990 (1989), and other promoters (e.g., cell promoters such as a eukaryotic cell promoter including, but not limited to, histone, RNA polymerase III, and β -actin promoter). Examples of other viral promoters that can be used include, but are not limited to, adenovirus promoter, thymidine kinase (TK) promoter, and B19 Parvovirus promoter. Persons skilled in the art can readily select a suitable promoter based on the teaching of the present invention.

[0058] A nucleic acid sequence that encodes the polypeptide of the present invention is under the control of a suitable promoter. Examples of suitable promoters that can be used include, but are not limited to, adenovirus promoter such as adenovirus major late promoter, heterologous promoter such as CMV promoter, respiratory syncytial virus (RSV) promoter, inducible promoter such as MMT promoter or metallothionein promoter, heat shock promoter, albumin promoter, ApoA1 promoter, human globin promoter, viral thymidine kinase promoter such as herpes simplex thymidine kinase promoter, retrovirus LTR including the aforementioned modified retrovirus LTR, β -actin promoter, and human growth hormone promoter. A promoter may be of a native type that controls the gene encoding polypeptides. A retrovirus plasmid vector is used to transduce the packaging cell line to form a producer cell line.

[0059] Examples of packaging cells to be transfected include, but are not limited to, PE501, PA317, Y-2, Y-AM, PA12, T19-14X, VT-19-17-H2, YCRE, YCRIP, GP+E-86, GP+envAm12, and the DAN cell line described in Miller, *Human Gene Therapy* 1: pp. 5-14 (1990).

[0060] A vector is transduced in a packaging cell by a means known in the art. Examples of such means include, but are not limited to, electroporation, the use of a liposome, and CaPO_4 precipitation. Alternatively, a retrovirus plasmid vector is sealed in a liposome or bound to a lipid to be administered to a host. A producer cell line produces infectious retrovirus vector particles comprising nucleic acid sequences encoding polypeptides. Such retrovirus vector particles are used to transduce eukaryotic cells *in vitro* or *in vivo*.

[0061] The transduced eukaryotic cells express nucleic acid sequences encoding polypeptides. Examples of eukaryotic cells that may be transduced include, but are not limited to, germinal stem cells, embryonal carcinoma cells, hematopoietic stem cells, hepatic cells, fibroblasts, sarcomas, keratinocytes, endothelial cells, and bronchial epithelial cells.

[0062] The protease of the present invention, an antibody against this protease, an antagonist of this protease, an inhibitor, an agonist, an activity modifier, or the like can be diluted with physiological saline, buffer, or the like to prepare a formulation. Thus, a pharmaceutical composition can be obtained. The pH value of the formulation is preferably between acidulous and neutral: close to the pH level of body fluid. The lower limit thereof is preferably between 5.0 and 6.4, and the upper limit is preferably between 6.4 and 7.4. Alternatively, the formulation can be provided in a state that allows storage for a long period of time, e.g., in a lyophilized state. In such a case, the formulation can be used by being dissolved in water, physiological saline, buffer, or the like at a desired concentration level at the time of use.

[0063] The formulation of the present invention may comprise a pharmacologically acceptable additive, such as a carrier, excipient, or diluent that is commonly used for pharmaceuticals, a stabilizer, or pharmaceutically necessary ingredients. Examples of a stabilizer include monosaccharides such as glucose, disaccharides such as saccharose and maltose, sugar alcohols such as mannitol and sorbitol, neutral salts such as sodium chloride, amino acids such as glycine, nonionic surfactants such as polyethylene glycol, polyoxyethylene and polyoxypropylene copolymers (Pluronic), polyoxyethylene sorbitan fatty acid ester (Tween), and human albumin. Addition thereof in amounts of about 1 to 10 w/v% is preferable.

[0064] An effective amount of the pharmaceutical composition of the present invention can be administered by, for example, intravenous injection, intramuscular injection, or hypodermic injection in one or several separate dosages. The dosage varies depending on symptom, age, body weight, or other factors, and it is preferably 0.001 mg to 100 mg per dose.

[0065] Also, sense or antisense DNA encoding the protease of the present invention can be similarly prepared in a formulation to obtain a pharmaceutical composition.

[0066] Further, the present invention includes methods for inhibiting platelet plug formation involved with heart infarction or brain infarction, methods for inhibiting arteriosclerosis, methods for preventing restenosis, reembo- lization, or infarction involved with PTCA, methods for preventing reembo- lization involved with PTCR, and methods for preventing platelet plug formation caused by HUS or O-157 through the administration of the peptide, protein, and DNA of the present invention. Furthermore, the present invention includes the use of the peptide, protein, and DNA of the present invention in the production of pharmaceuticals for inhibiting platelet plug formation involved with heart infarction or brain infarction, pharmaceuticals for inhibiting arteriosclerosis, pharmaceuticals for preventing restenosis, reembo-

ization, or infarction involved with PTCA, pharmaceuticals for preventing reemboilization involved with PTCR, and pharmaceuticals for preventing platelet plug formation caused by HUS or O-157.

[0067] The peptide or protein of the present invention is used as a leading substance for amino acid modification. This enables the preparation of a molecule having activity that is different from that of the protease of the present invention. An example thereof is a variant molecule that can be obtained by preparing an antagonist, which is obtained by preparing a variant deactivated through amino acid substitution between an amino acid residue located around the active center in the metalloprotease domain and another amino acid, separating a molecule recognition site from a catalytic site, or varying one or both of these sites.

[0068] The use of an evaluation system for the vWF-cleaving activity described herein enables the production of an antagonist/agonist. For example, an effective antagonist can be a small organic molecule, a peptide, or a polypeptide. An example thereof is an antibody that is bound to the polypeptide of the present invention, thereby inhibiting or eliminating its activity.

[0069] Similarly, the use of the aforementioned evaluation system for vWF-cleaving activity enables the screening for a compound that is capable of cleaving vWF. In such a case, the cleaving activity of the test compound may be evaluated using the aforementioned evaluation system.

Brief Description of the Drawings

[0070]

Fig. 1 is a diagram showing the vWF multimer structure and the point cleaved by the vWF-cleaving protease.

Fig. 2 is a photograph showing the result of vWF multimer analysis (agarose electrophoresis).

Fig. 3 is a photograph showing the result of SDS-PAGE (5% gel) for analyzing the vWF-cleaving activity of each plasma fraction under reducing conditions.

Fig. 4 is a photograph showing the result of SDS-PAGE (5% gel) for analyzing the solubilized sample of fraction 1 (F1) paste under non-reducing conditions.

Fig. 5 is a photograph showing the result of analyzing vWF-cleaving protease fractions after being subjected to gel filtration chromatography three times using the solubilized sample of F1 paste as a starting material. Fig. 5A is a chart showing gel filtration chromatography, Fig. 5B shows the result of SDS-PAGE on fractions under non-reducing conditions, and Fig. 5C shows the results of SDS-PAGE on vWF-cleaving activity under reducing conditions.

Fig. 6 is a photograph showing the results of analyzing vWF-cleaving protease fractions in which the fraction collected by gel filtration chromatography is purified by DEAE anion exchange chromatography. Fig. 6A is a chart showing gel filtration chromatography, Fig. 6B shows the result of SDS-PAGE (8% gel) on elution fractions under non-reducing conditions, and Fig. 6C shows the results of SDS-PAGE on vWF-cleaving activity under reducing conditions. In Fig. 6C, three bands indicate an intact vWF molecule (remaining uncleaved), a vWF cleavage fragment, and a vWF cleavage fragment, respectively, as in Fig. 5C.

Fig. 7 is a photograph showing an electrophoresed fragment obtained when the vWF-cleaving protease fraction purified and concentrated by DEAE anion exchange chromatography is further purified by Biophoresis-based SDS-PAGE (non-reducing conditions).

Fig. 8 is a photograph showing the result of electrophoresis on a fraction obtained by further purifying a vWF-cleaving protease fraction by Biophoresis-based SDS-PAGE for analyzing vWF-cleaving protease activity and SDS-PAGE on active fractions under reducing conditions. Fig. 8A shows the results of SDS-PAGE for analyzing vWF-cleaving protease activity under non-reducing conditions, and Fig. 8B shows the results of SDS-PAGE for analyzing active fractions under reducing conditions.

Fig. 9 relates to the identification of the vWF-cleaving protease gene, which is a diagram showing primers used for amplifying the gene fragment for a Northern blot probe.

Fig. 10 relates to the identification of the vWF-cleaving protease gene, which is a photograph showing Northern blot autoradiography. Fig. 10A shows the results obtained when the protease-encoding gene is used as a probe, and Fig. 10B shows the results obtained when a β -actin probe (RNA control) is used.

Fig. 11 relates to the identification of the vWF-cleaving protease gene, and is a diagram showing the locations and the sequences of the primers used in the RACE experiments.

Fig. 12 is a diagram showing the locations of primers designed for cloning full-length cDNA.

Fig. 13 is a diagram showing a process for constructing a vector containing full-length cDNA.

Fig. 14 is a photograph showing the expression in various cell lines (Western blotting under reducing conditions using anti-FLAG antibody, where the mock is prepared by inversely inserting a gene in an expression vector). In Fig. 14, each lane shows the results using the indicated sample.

Lane 1: Mock (host: 293 cell)
 Lane 2: vWF-cleaving protease, cDNA+FLAG (host: 293 cell)
 Lane 3: Mock (host: HepG2 cell)
 Lane 4: vWF-cleaving protease, cDNA+FLAG (host: HepG2 cell)
 Lane 5: Mock (host: Hela cell)
 Lane 6: vWF-cleaving protease, cDNA+FLAG (host: Hela cell)

Fig. 15 is a photograph showing the activity assay of recombinant expression protease (analysis of vWF-cleavage by SDS-PAGE under non-reducing conditions, where the mock is prepared by inversely inserting a gene in an expression vector). In Fig. 15, each lane shows the results using the indicated sample.

Lane 1: Mock (host: Hela cell)
 Lane 2: Supernatant in which vWF-cleaving protease was expressed (host: Hela cell)
 Lane 3: Mock (host: HepG2 cell)
 Lane 4: Supernatant in which vWF-cleaving protease was expressed (host: HepG2 cell)
 Lane 5: Mock (host: 293 cell)
 Lane 6: Supernatant in which vWF-cleaving protease was expressed (host: 293 cell)
 Lane 7: Mock (host: BHK cell)
 Lane 8: Supernatant in which vWF-cleaving protease was expressed (host: BHK cell)
 Lane 9: Mock (host: COS cell)
 Lane 10: Supernatant in which vWF-cleaving protease was expressed (host: COS cell)
 Lane 11: Mock (host: CHO cell)
 Lane 12: Supernatant in which vWF-cleaving protease was expressed (host: CHO cell)

Fig. 16 is a photograph showing the result of Western blotting using an antibody established against the protease of the present invention, wherein Western blotting is carried out for various antisera using the 293 cell as a host and a recombinant vWF-cleaving protease. In Fig. 16, each lane shows the results obtained with the use of the indicated sample.

Lane 1: Mouse antiserum (prepared by administering purified protein)
 Lane 2: Rabbit antiserum (prepared by hypodermically administering an expression vector to a rabbit)
 Lane 3: Untreated rabbit antiserum
 Lane 4: Rabbit antiserum (prepared by administering KLH-conjugated partial synthetic peptide)

Fig. 17 is a photograph showing the result of Western blotting using an antibody established against the protease of the present invention, wherein various samples derived from human plasma and recombinant expression units are detected using rabbit antiserum obtained by administering full-length cDNA, of vWF-cleaving protease. In Fig. 17, each lane shows the results obtained with the use of the indicated sample.

Lane 1: Partially purified sample derived from human plasma cryoprecipitate
 Lane 2: Purified vWF-cleaving protease derived from human plasma
 Lane 3: Gel-filtrated FI paste sample obtained from pooled human plasma
 Lane 4: Recombinant vWF-cleaving protease (host: 293 cell)
 Lane 5: Recombinant vWF-cleaving protease (host: Hela cell)

Fig. 18 is a photograph showing the result of Western blotting using an antibody established against the protease of the present invention, wherein rabbit antiserum obtained by immunizing a rabbit with a partially synthesized peptide of the vWF-cleaving protease is used to confirm the vWF-cleaving protease in healthy human plasma and that in the plasma and gene recombinant vWF-cleaving protease of a TTP patient. In Fig. 18, each lane shows the results obtained with the use of the indicated sample.

Lane 1: Gel-filtrated FI paste sample obtained from pooled human plasma
 Lane 2: Normal human plasma 1
 Lane 3: Normal human plasma 2
 Lane 4: Normal human plasma 3
 Lane 5: TTP patient's plasma 1
 Lane 6: TTP patient's plasma 2
 Lane 7: Recombinant vWF-cleaving protease (host: 293 cell)

Lane 8: Recombinant vWF-cleaving protease (host: Hela cell)

Fig. 19 is a diagram showing the result of ELISA using an antibody prepared against the vWF-cleaving protease. Fig. 20 is a photograph showing the result of SDS-PAGE (silver staining) analyzing each fraction of affinity purified vWF-cleaving protease using an antibody under reducing conditions. In Fig. 20, each lane shows the results obtained with the use of the indicated sample.

Lane 1: Applied culture supernatant (diluted 10-fold)

Lane 2: Passed-through fraction

Lane 3: Washed fraction

Lane 4: Elution fraction

Fig. 21 is a photograph showing the results of evaluating neutralizing activity using an antibody (SDS-PAGE for analyzing vWF-cleaving activity under non-reducing conditions). In Fig. 21, each lane shows the results obtained with the use of the indicated sample.

Lane 1: vWF-cleaving protease solution: normal rabbit serum = 1:1

Lane 2: vWF-cleaving protease solution: normal rabbit serum (diluted 5-fold) = 1:1

Lane 3: vWF-cleaving protease solution: peptide-immunized rabbit serum = 1:1

Lane 4: vWF-cleaving protease solution: peptide-immunized rabbit serum (diluted 5-fold) = 1:1

Lane 5: vWF-cleaving protease solution: recombinant protein-immunized rabbit serum = 1:1

Lane 6: vWF-cleaving protease solution: recombinant protein-immunized rabbit serum (diluted 5-fold) = 1:1

Lane 7: vWF-cleaving protease solution: 10mM EDTA = 1:1

Lane 8: vWF-cleaving protease solution: buffer only = 1:1

Lane 9: buffer (without vWF-cleaving protease) : buffer = 1:1

Fig. 22 is a diagram showing the construction of an expression vector for a molecular species lacking a C-terminal domain.

Best Modes for Carrying out the Invention

[0071] The present invention is hereafter described in detail with reference to the following examples, although it is not limited to these examples.

Example 1

(Preparation of vWF)

[0072] A plasma cryoprecipitation (2 g) was dissolved in 20 ml of buffer (0.01 % Tween-80/50 mM Tris-HCl/100 mM NaCl, pH 7.4), and the resultant was subjected to gel filtration using a Sephacryl S-500 HR Column (2.6 x 90 cm, Amersham Pharmacia) to prepare vWF. Fractions were recovered at a flow rate of 2 ml/min in amounts of 6 ml each. vWF was analyzed by Western blotting using a peroxidase-labeled rabbit anti-human vWF antibody (DAKO), and high-molecular-weight vWF fractions were pooled. The pooled fractions were subjected to multimer analysis using agarose electrophoresis as described below.

[0073] As shown in Fig. 1, vWF originally has a multimer structure in which vWF monomer molecules are polymerized with each other at their N-terminuses or at their C-terminuses, and vWF is subjected to partial hydrolysis by the vWF-specific cleaving protease. As a result of the analysis, as shown in Fig. 2, the purified vWF exhibited a multimer pattern based on agarose electrophoresis approximately equivalent to that in the plasma of a healthy person (the ladder in the drawing shows the electrophoresis pattern of vWF having a multimer structure, and the upper portion indicates vWF with advanced polymerization). This can prepare vWF comprising substantially no impurities that degrade it, and this fraction was used as a substrate when assaying the vWF-cleaving activity as described below.

Example 2

(vWF-cleaving reaction)

[0074] vWF-cleaving activity was assayed as follows. A sample comprising 10 mM barium chloride (final concentration) was pre-incubated at 37°C for 5 minutes to activate protease. A buffer (15 to 20 ml, 1.5 M urea/5 mM Tris-HCl,

pH 8.0) was placed in a 50 ml Falcon Tube. Subsequently, a membrane filter (0.025 μ m, Millipore) was floated therein, and 100 μ l of activated sample prepared by mixing with 50 μ l of vWF substrate solution was added. The resultant was allowed to stand in an incubator (37°C) overnight and recovered from the filter on the next day. The recovered sample was evaluated based on the vWF cleavage pattern as described below in the "SDS-PAGE" section.

SDS-PAGE

[0075] SDS-5% polyacrylamide gel was autologously prepared and used. An SDS electrophoresis buffer (2 μ l, in the presence or absence of a reducing agent, i.e., 2-mercaptoethanol) was added to 10 μ l of the sample described in the "vWF-cleaving activity assay" section, and the resultant was boiled for 3 minutes to prepare an electrophoresis sample. The gel was subjected to electrophoresis at 30 mA for 1 hour and then stained with the Gel Code Blue Stain Reagent (PIERCE) utilizing CBB staining. As shown in Fig. 1, activity is evaluated based on the development of a cleavage fragment and the presence or absence of fragments remaining uncleaved under reducing or non-reducing conditions. This is more specifically described in Example 3 and Fig. 3 below.

Multimer analysis utilizing agarose electrophoresis

Preparation of gel, electrophoresis

[0076] Low gelling temperature agarose (Type VII, Sigma) was added to 375 mM Tris-HCl (pH 6.8) until a concentration of 1.4% was reached, followed by heating in a microwave oven to completely dissolve the gel. Thereafter, 0.1% SDS was added, and the resultant was maintained at 56°C. The resultant was made to flow into a gel mold and solidified by cooling at 4°C overnight (running gel). The next day, high gelling temperature agarose (SeaKem) was mixed with 375 mM Tris-HCl (pH 6.8) until a concentration of 0.8% was reached, and dissolved by boiling in a microwave oven. Thereafter, the resultant was maintained at 56°C (stacking gel). The gel prepared on the previous day was cleaved, leaving a 10-cm fraction from the end uncleaved. The aforementioned gel was made to flow into the cleaved portion, and the gel was made to keep flowing at 4°C for at least 3 hours, followed by solidification. Pyronin Y was added to the sample described in the "vWF cleaving activity assay" section above, and the gel was prepared under non-reducing conditions without boiling. The gel was subjected to electrophoresis at 10 mA for at least 24 hours using an SDS-PAGE buffer.

Western blotting

[0077] After the electrophoresis, the gel was immersed in a transcription buffer (0.005% SDS, 50 mM phosphate buffer, pH 7.4) for 10 minutes, and the resultant was transferred to a nitrocellulose membrane using a transcription apparatus at 4°C at 0.5 A overnight. Blocking was performed using a blotting solution (5% skim milk, PBS) for 30 minutes, and the gel was then allowed to react for at least 6 hours with the peroxidase-labeled rabbit anti-human vWF antibody (DAKO), which was diluted 1,000-fold with the blotting solution. Thereafter, the gel was washed three times with the blotting solution and once with PBS, and color was developed using Konica Immunostain HRP-1000 (Konica), which was a substrate reaction solution for peroxidase. The purified vWF analyzed in this assay was found to have been undegraded, but was sufficiently usable as a substrate in the present invention (Fig. 2).

Example 3

(Preparation of vWF-cleaving protease)

[0078] Plasma was subjected to ethanol fractionation developed by Cohn. A protease having high vWF-cleaving activity (one with high specific activity) when protein levels in four fractions (i.e., starting plasma, cryoprecipitate, fraction I (FI) supernatant, and a paste) are made equivalent to each other was selected. As shown in Fig. 3, the protease activity was highest in the FI paste. The N-terminal sequence of this cleavage fragment was analyzed, and as a result, activity derived from the cryoprecipitate and the FI paste were found to cleave the peptide bond between residues Tyr 842 and Met 843. Thus, the FI paste was determined to be a main starting material for purification thereafter.

Solubilization of FI paste

[0079] The FI paste was fractionated in fractions of 12 g each and then cryopreserved. The paste was allowed to melt at 4°C the day before its use. The next day, 120 ml of solubilizing buffer (0.05% azide, 50 mM Tris-HCl (pH 7.4), 100 mM NaCl) was added at 10 mg/ml, and the mixture was stirred at 37°C for 2 hours. The product was centrifuged at 10,000 rpm for 10 minutes, and the supernatant was then recovered, followed by filtration with a prefilter, a 5.0 μ m

filter, and a 0.8 μ m filter in that order. The resultant was determined to be a solubilized sample. Fig. 4 shows the result of SDS-PAGE of the solubilized sample.

Gel filtration chromatography of vWF-cleaving protease

[0080] The solubilized F1 paste was applied to a Sephacryl S-300 HR Column (5 x 90 cm, Amersham Pharmacia) to conduct the first gel filtration. A buffer comprising 0.05 % azide, 50 mM Tris-HCl (pH 7.4), and 100 mM NaCl (hereinafter referred to as an "elution buffer"), which was the same as the solubilizing buffer, was used. The flow rate was 5 ml/min, fractionation was initiated at 600 ml after the sample application, and fractions were recovered in amounts of 10 ml each. Fractions were subjected to the vWF-cleaving reaction, and their activities were then analyzed by SDS-PAGE. Fractions that exhibited protease activity were pooled, and a small amount of saturated ammonium sulfate was gradually added dropwise thereto until a final concentration of 33% saturation was reached. The mixture was further allowed to stand at 4°C overnight. The next day, the product was centrifuged at 10,000 rpm for 10 minutes, and an active fraction of interest was recovered as a precipitate. The procedures comprising solubilization, gel filtration, and ammonium sulfate precipitation were performed for 5 batches and the resultant was cryopreserved at -20°C.

[0081] The ammonium sulfate precipitates (2 to 3 batches) obtained by the first gel filtration were dissolved in 50 ml of elution buffer, and passed through the Sephacryl S-300 HR Column (5 x 90 cm) in the same manner as in the first gel filtration to perform the second gel filtration. The elution buffer, conditions, operations, and the like were the same as those in the first gel filtration. Fractions were subjected to the vWF-cleaving reaction, and their activities were then analyzed by SDS-PAGE. Fractions with activity were pooled, and ammonium sulfate precipitation was similarly performed. These procedures were repeated two times.

[0082] The ammonium sulfate precipitates (2 batches) obtained by the second gel filtration were dissolved in 50 ml of elution buffer, and applied to the Sephacryl S-300 HR Column (5 x 90 cm) in the same manner as in the first and the second gel filtration to perform the third gel filtration. The elution buffer, conditions, operations, and the like were the same as those in the first and the second gel filtration. Fractions were subjected to the vWF-cleaving reaction, and their activities were then analyzed by SDS-PAGE, followed by pooling. Fig. 5 shows SDS-PAGE for analyzing these fractions and that for analyzing vWF-cleaving activity. Based on the patterns of gel filtration and the data showing activity, the protease of the present invention was found to be eluted in the region between fraction 37 and fraction 47. Based on a separately conducted elution experiment for high-molecular-weight gel filtration marker (Amersham Pharmacia), this site of elution was deduced to have a molecular weight equivalent to 150 to 300 kDa. In this phase, considerable amounts of impurities were still present.

DEAE anion exchange chromatography

[0083] The pooled fraction obtained by three gel filtration operations was subjected to dialysis overnight with a buffer comprising 50 mM Tris-HCl and 50 mM NaCl (pH 7.1). After the dialysis, anion exchange chromatography was performed using a 5 ml HiTrap DEAE-Sepharose Fast Flow Column (Pharmacia) to conduct further purification and concentration. Equilibrating and washing were performed using a buffer comprising 50 mM Tris-HCl (pH 7.1), and elution was performed using 0.25 M NaCl. The flow rate was 5 ml/min, and 5 fractions of 5 ml each were recovered and pooled. Fig. 6 shows the results of SDS-PAGE for analyzing elution fractions and those for analyzing vWF-cleaving activity. Based on SDS-PAGE for activity assay, the protease of the present invention having vWF-cleaving activity was considerably effectively concentrated in the elution fraction.

Fractionation utilizing SDS-PAGE

[0084] The sample (5 ml) purified and concentrated by DEAE anion exchange chromatography was further concentrated to 0.5 ml using Centricon (molecular weight cut off: 10,000 Da, Amicon). The protease of the present invention was isolated by Biophoresis III (Atto Corporation) utilizing SDS-PAGE. In accordance with the Laemmli method (Nature, vol. 227, 680-685, 1970), a buffer for electrophoresis tanks was prepared, and developed with 8% polyacrylamide gel to recover the electrophoresis fraction. Fig. 7 shows the result of SDS-PAGE for analyzing the recovered fractions. The buffer used for recovery was comprised of 50 mM Tris-HCl and 10% glycerol (pH 8.8). As is apparent from Fig. 7, this process according to the present invention has a high ability to produce separation. Fig. 8 shows the results of analyzing activity of a fraction further purified by electrophoresis and the results of SDS-PAGE for analyzing active fractions. The protease of the present invention can be recovered as an active molecule even after SDS-PAGE. When the activity of this protease in the plasma is determined to be 1 in terms of specific activity, a degree of purification of 30,000- to 100,000-fold was deduced to be achieved based on the average protein content in the plasma (60 mg/ml).

Example 4

(Partial amino acid sequencing)

[0085] The partial amino acid sequence of the isolated protease was determined. This protease, which was isolated using Biophoresis, was transferred to a PVDF membrane after SDS-PAGE by a conventional technique, air-dried, and then subjected to analysis using the automated protein sequencer (model 492; PE Applied Biosystems). As a result, the vWF-cleaving protease of the present invention isolated under the above conditions was found to comprise a polypeptide chain having a molecular weight of 105 to 160 kDa in SDS-PAGE under reducing conditions. This protease was also found to have, as a partial sequence, Leu-Leu-Val-Ala-Val, and preferably Ala-Ala-Gly-Gly-Ile-Leu-His-Leu-Glu-Leu-Leu-Val-Ala-Val.

Deduction of isolated protease utilizing bioinformatics

[0086] At present, bioinformatics enables the deduction of full nucleotide sequences encoding a polypeptide without substantial gene cloning through collation with information in the database accumulated in the past (BIOINFORMATICS: A Practical Guide to the Analysis of Genes and Proteins, edited by Andreas D. Baxeavanis and B. F. Francis Ouellette). Based on the partial amino acid sequencing by the aforementioned process (Ala-Ala-Gly-Gly-Ile-Leu-His-Leu-Glu-Leu-Leu-Val-Ala-Val), the database was searched by the tblastn program. As a result, a chromosome clone (AL158826) that was deduced to encode the protease of the present invention was identified by genomic database search. Further, a part of the protease of interest as the expressed sequence tag (EST) and a clone that was deduced to be a part of the polypeptide encoded by the aforementioned genome (AI346761 and AJ011374) were identified. The amino acid sequence as shown in SEQ ID NO: 3 or 7 was deduced based thereon to be an active vWF-cleaving protease site.

Example 5

(Gene identification)

[0087] Synthesis of all the following synthetic primers was performed by Greiner Japan Co.Ltd. by request. Further, reagents used for gene recombination were those manufactured by TAKARA, TOYOBO, and New England Biolabs unless otherwise specified.

Preparation of a gene fragment as a Northern blotting probe

[0088] A sense primer (SEQ ID NO: 9) and an antisense primer (SEQ ID NO: 10) were prepared. PCR was carried out using Universal QUICK-Clone™ cDNA (Clontech), which was a mixture of cDNA derived from normal human tissue, as a template and TaKaRa LA Taq with GC rich buffer. A gene sandwiched between these primers was amplified, and the amplified fragment was cloned using a TOPO TA cloning™ kit (Invitrogen). DNAs having the nucleotide sequence as shown in SEQ ID NO: 6 were isolated from several clones.

[0089] A vector portion was removed from this cloned DNA by EcoRI digestion, separated and purified by agarose electrophoresis, and the resultant was determined to be a template for preparing probes for Northern blotting.

Northern blotting

[0090] The gene fragment prepared above was employed as a template to prepare a radioactive probe using [α -³²P] dCTP (Amersham Pharmacia) and a BcaBEST™ labeling kit (TAKARA). Hybridization was carried out using the Human 12-lane Multiple Tissue Northern Blots™ (Clontech) filter in accordance with the method described in Molecular Cloning 2nd Edition, pp. 9.52-9.55. Detection was carried out by autoradiography. As shown in Fig. 10, mRNA encoding the protease was expressed mainly in the liver. The size of this mRNA was found to be more than 4.4 kb.

Isolation and identification of gene encoding the protease

[0091] As a result of Northern blotting, mRNA was found to be expressed mainly in the liver. Thus, the protease gene of the present invention was isolated and identified in accordance with the RACE technique using normal human liver-derived poly A⁺ RNA and Marathon-Ready™ cDNA (Clontech).

[0092] More specifically, the first PCR was carried out as 5' RACE using normal human liver-derived Marathon-Ready™ cDNA in accordance with the product's manual and using the AP-1 primer attached to the kit and antisense

primers (SEQ ID NOs: 11 to 13) arbitrarily selected from the group of Gene Specific Primers (GSP) excluding the primer 1 located in the uppermost stream as shown in Fig. 11. Nested PCR (the second PCR) was then carried out using the AP-2 primer located in the inside thereof and the antisense primer located in the inside of the primer used for the first PCR as shown in Fig. 11. Thereafter, TA cloning was carried out. Genes were prepared from the developed colonies in accordance with a conventional technique (Molecular Cloning 2nd Edition, pp. 1.25-1.28), and nucleic acid sequences were decoded using an automatic DNA sequencer. The primer used for sequencing was the primer used for PCR or a primer located in the inside thereof. Further, the primer was designed based on the sequence determined after serial decoding.

[0093] 3' RACE was started from normal human liver-derived poly A⁺ RNA using the 3'-Full RACE Core Set (TAKARA), and reverse transcription was carried out in accordance with the attached manual using the attached oligo dT primer. The band amplified by PCR using the sense primer (SEQ ID NO:14) located at "primer 2" in Fig. 11 and the attached oligo dT primer was separated by agarose electrophoresis and extracted, followed by TA cloning. Genes were prepared from the developed colonies, and nucleic acid sequences were decoded using an automatic DNA sequencer. A primer used for sequencing was designed based on the sequence determined after serial decoding.

Example 6

(Preparation of a vector comprising full-length cDNA 1)

[0094] cDNA encoding the protein was subjected to one-stage PCR by, for example, using a sense primer 1 (SEQ ID NO: 22) comprising an XhoI restriction site and an initiation codon and an antisense primer 2 (SEQ ID NO: 23) comprising an Sall restriction site and a termination codon (see Fig. 12), using the aforementioned normal human liver-derived Marathon-ReadyTM cDNA as a template and the TaKaRa LA Taq with GC rich buffer, followed by the aforementioned TA cloning. Thereafter, the full length of the product was confirmed using an automatic DNA sequencer.

Example 7

(Preparation of a vector comprising full-length cDNA 2)

[0095] Restriction sites Accl and AvrII that cleaved cDNA only at one point on the inner sequence of the cDNA (SEQ ID NO: 15) encoding the protein were found. With the use thereof, full-length cDNA was divided into three fragments as shown in Fig. 12. A fragment 1 sandwiched between the sense primer 1 (SEQ ID NO: 22) and the antisense primer 3 (SEQ ID NO: 24), a fragment 2 sandwiched between the sense primer 4 (SEQ ID NO: 25) and the antisense primer 5 (SEQ ID NO: 26), and a fragment 3 sandwiched between the sense primer 6 (SEQ ID NO: 27) and the antisense primer 2 (SEQ ID NO: 23) were provided, respectively, in each of the above three fragments. Each fragment was subjected to PCR using the aforementioned normal human liver-derived Marathon-ReadyTM cDNA as a template and TaKaRa LA Taq with GC rich buffer, followed by the aforementioned TA cloning. The full length of the product was confirmed using an automatic DNA sequencer. Further, the pCR 2.1 vector included in the aforementioned TA cloning kit was subjected to self ligation, the ligation product was cleaved with XhoI/HindIII, ligated to a linker comprising XhoI/AccI/AvrII/HindIII (prepared by annealing the synthetic DNA as shown in SEQ ID NO: 28 or 29), and the three aforementioned fragments were sequentially ligated in a conventional manner to bind them. Thus, cDNA comprising the entire region was prepared (see Fig. 13).

Example 8

(Preparation of an expression vector comprising full-length cDNA: an animal cell host)

[0096] DNA obtained in Example 6 or 7 was digested with restriction enzymes XhoI/Sall, ligated to, for example the Sall site in the pCAG vector (Niwa, H. et al., Gene, vol. 108, 193-199), and the direction of the insertion and the full-length sequence were confirmed using an automatic DNA sequencer.

Example 9

(Transfection of an expression vector comprising full-length cDNA into an animal cell)

[0097] The animal cell expression vector prepared in Example 8 was transfected in the following manner using the 293 cell (human embryonic kidney cell line), the Hela cell, and the HepG2 cell. At the outset, cells were disseminated at 1 to 3 x 10⁵ cells per 35 mm dish 24 hours before the transfection. The next day, 2 µl of polyamine transfection

reagent, TransIT (TAKARA), per μg of the expression vector, were added to 100 μl of a serum-free medium such as Opti-MEM to prepare a complex with DNA in accordance with the instructions included with the reagent. Thereafter, the complex was added dropwise to the various types of previously prepared cells, and the resultants were incubated for 2 to 8 hours, followed by medium exchange. The medium was further exchanged three days later with the selective medium to which G418 had been added. Thereafter, medium was exchanged every three days to produce a stably expressed strain. An example thereof is shown in Fig. 14 as a temporarily expressed strain comprising an FLAG epitope tag at its C-terminus. Detection was carried out by Western blotting using the anti-FLAG-M2 antibody (Kodack) and staining with anti-mouse Ig-alkaline phosphatase-labeled antibody system. The recombinant strain expressed using cDNA as shown in this example exhibited a molecular size of about 250 kDa under reducing conditions. This molecular size was also found in the plasma of a healthy human (Fig. 18, Example 14 below). Several different molecular species of this protease are found to be present in the human plasma, which could be caused by the presence of the alternative splicing products (SEQ ID NOS: 6 to 21) observed at the time of gene cloning, difference in post-translational modification such as sugar chain addition, or degradation during purification (described in Example 14 and in Fig. 17 of the present invention and Gerritsen et al., Blood, vol. 98, 1654-1661 (2001)).

[0098] Subsequently, the vWF-cleaving activity of the recombinant strain was confirmed by the method described in Example 2 (Fig. 15). As a result, the human plasma-derived protease and the gene recombinant product of the present invention were found to exhibit the same vWF-cleaving activities.

Example 10

(Preparation of an expression vector comprising partial cDNA: an *E. coli* host)

[0099] Partial cDNA encoding the metalloprotease domain of the protein was subjected to PCR using a sense primer comprising an NcoI restriction site and an initiation codon (SEQ ID NO: 30), and an antisense primer comprising an HindIII restriction site and a termination codon (SEQ ID NO: 31), the aforementioned normal human liver-derived Marathon-Ready™ cDNA or the cDNA obtained in Example 6 or 7 as a template, and the TaKaRa LA Taq with GC rich buffer. The PCR product was then digested with NcoI/HindIII, ligated to the NcoI/HindIII digest of an *E. coli* expression vector such as pUT1 (Soejima et al., J. Biochem. Tokyo, vol. 130, 269-277 (2001)), and transformed to the *E. coli* competent cell JM 109 by a conventional technique. Several clones were collected from the formed colony group, and genes were prepared therefrom. Thereafter, the resulting genes were confirmed to be the genes encoding the polypeptide, wherein the nucleic acid sequence of the insertion site of the plasmid vector was equivalent to SEQ ID NO: 32 or substantially represented by SEQ ID NO: 33, using an automatic DNA sequencer.

Example 11

(Expression of partial cDNA-containing expression vector in *E. coli*)

[0100] An *E. coli* host with the expression vector constructed in Example 10 introduced therein was precultured in 200 ml of LB medium comprising 50 $\mu\text{g}/\text{ml}$ ampicillin at 30°C overnight. The resultant was sowed in a fermenter comprising 8 liters of LB medium, and culture was conducted at 30°C until the turbidity at 600 nm became 0.2 to 0.5. Thereafter, isopropyl-1-thio- β -D-galactopyranoside was added to a final concentration of 1 mM, and the mixture was further cultured overnight to induce the metalloprotease domain of the protein to be expressed. The cultured *E. coli* were collected using a centrifuge (4°C for 30 minutes).

[0101] Subsequently, the collected *E. coli* pellet was resuspended in distilled water, and lysozyme (final concentration: 0.6 mg/ml) was added thereto. The mixture was stirred at room temperature for 30 minutes, allowed to stand at 4°C overnight, and cells were then destroyed. After the ultrasonication, centrifugation was carried out using a centrifuge (4°C for 20 minutes), and the pellet was recovered. The recovered pellet was resuspended in a buffer comprising 50 mM Tris, 10 mM EDTA, and 1% Triton X-100 (pH 8.0). These procedures of centrifugation, ultrasonication, and resuspension were repeated several times, and the pellet was then resuspended in distilled water. Similarly, procedures of centrifugation, ultrasonication, and resuspension were repeated several times to recover an inclusion body. This inclusion body was used as an antigen when producing an antibody.

Example 12

(Isolation of homologous gene of other animal species)

[0102] The nucleic acid sequence as shown in SEQ ID NO: 15 was used as a probe, and a homology search was conducted using the BLASTN program at the GenomeNet WWW server (<http://www.genome.ad.jp/>). As a result, chro-

mosome clones AC091762 and AC090008 that were mapped at mouse chromosome 10 were obtained. Based on these sequences, a mouse homolog of the protease of the present invention as shown in SEQ ID NO: 34 was deduced. A new primer was designed from this sequence, and Northern blot analysis was conducted by the technique used in isolating and identifying the gene encoding the human vWF-cleaving protease. Thus, the occurrence of the specific expression in the liver was observed as with the case of humans. Further, normal mouse liver-derived poly A+ RNA and Marathon-Ready™ cDNA (Clontech) were used to isolate and identify the protease gene of the present invention by the RACE technique as in the case of humans. As a result, the mouse homologous gene sequences of the protease as shown in SEQ ID NOs: 35 and 36 were determined.

[0103] Based on the thus determined mouse homologous partial sequence, the Exon/Intron structure on the 5' side of the aforementioned mouse chromosome 10 was determined. In accordance with a conventional technique (e.g., Gene Targeting: A Practical Approach First Edition, edited by A. L. Joyner, Teratocarcinomas and embryonic stem cell a practical approach), a targeting vector for knock-out (knock-in) mice can be prepared based thereon. This enabled the production of mutated mice. Further, this protein can be subjected to recombinant expression by a conventional technique.

Example 13

(Production of an antibody and construction of a detection system for the present protease using the antibody)

[0104] In accordance with a conventional technique (e.g., Current Protocols in Molecular Biology: Chapter 11 immunology, Antibody Engineering: A PRACTICAL APPROACH, edited by J. McCafferty et al. or ANTIBODY ENGINEERING second edition, edited by Carl A. K. Borrebaeck), an expression vector was administered to a mouse or rat. This expression vector comprises a substance prepared by optionally binding an antigen protein partially purified from human plasma or a synthetic peptide having a partial amino acid sequence thereof (e.g., a C-terminal peptide sequence (SEQ ID NO: 37) Phe-Ser-Pro-Ala-Pro-Gln-Pro-Arg-Arg-Leu-Leu-Pro-Gly-Pro-Gln-Glu-Asn-Ser-Val-Gln-Ser-Ser, which was one isoform of the protease of the present invention) to an optimal carrier substance such as KLH (Cys was added to, for example, the N- or C-terminus to facilitate KLH addition), the aforementioned gene recombinant protein, or a gene encoding this protein. Thus, a monoclonal antibody-expressing hybridoma was established, and a polyclonal antibody (antiserum) was produced.

[0105] Subsequently, the antibodies prepared by the various aforementioned techniques were used to detect the protease of the present invention by Western blotting in accordance with a conventional technique (e.g., Current Protocols in Molecular Biology: Chapter 10 analysis of proteins, Chapter 11 immunology). More specifically, the culture supernatant of the recombinant unit-expressing 293 cell obtained in the procedure as described in Example 9 was subjected to SDS-PAGE under non-reducing conditions, transferred to a PVDF membrane, and confirmed using mouse or rabbit antiserum to confirm the expression of the genetically recombinant unit (Fig. 16). As a result, a band that was deduced to be derived from the protease of the present invention was found in a molecular size range of 160 to 250 kDa. Subsequently, the protease of the present invention was detected using starting plasma or the like and a recombinant unit under non-reducing conditions. As a result, a band was found in 105 to 160 kDa or 160 to 250 kDa (Fig. 17). Also, a band derived from a similar recombinant unit was detected in a monoclonal antibody established by immunizing a recombinant protein (clone No. CPHSWH-10).

[0106] Further, the C-terminal peptide sequence Phe-Ser-Pro-Ala-Pro-Gln-Pro-Arg-Arg-Leu-Leu-Pro-Gly-Pro-Gln-Glu-Asn-Ser-Val-Gln-Ser-Ser (SEQ ID NO: 37), which was one isoform of the protease of the present invention, was bound to KLH. The resultant was used as an immunogen to obtain a peptide antibody. With the use thereof, the protease of the present invention was detected from the plasma of healthy persons, plasma of TTP patients, or a culture supernatant of the recombinant unit under reducing conditions. As a result, a band of approximately 250 kDa that was deduced to be a signal derived from the protease of the present invention was found, although it was not clear based on plasma derived from some TTP patients (Fig. 18).

[0107] Furthermore, enzyme immunoassay (ELISA) constructed by combining the obtained antibodies enabled the preparation of a calibration curve that is concentration-dependent at the culture supernatant level of the recombinant protein (Fig. 19). An example of ELISA is as follows. The obtained mouse anti-vWF-cleaving protease antibody was immobilized on the Maxisorp plate (Nunc), and 1/1, 1/2, and 1/4 diluents of the culture supernatant of the vWF-cleaving protease-temporarily expressing 293 cells were allowed to react in amounts of 100 µl/well (Mock supernatant as "0"). The plate was subjected to reaction, for example, at 37°C for 1 hour, and then washed with 0.05% Tween 20/TBS. Thereafter, the 100-fold diluted rabbit anti-vWF-cleaving protease antibody was allowed to react in amounts of 100 µl/well, for example, at 37°C for 1 hour, and the plate was washed with 0.05% Tween 20/TBS. The 1,000-fold diluted peroxidase-labeled anti-rabbit Ig antibody (BioRad) was then allowed to react in amounts of 100 µl/well, for example, at 37°C for 1 hour, and the plate was washed with 0.05% Tween 20/TBS. Thereafter, color was developed for a given period of time using a coloring substrate TMBZ, the reaction was terminated using 1M sulfuric acid as a termination

liquid, and the absorbance at 450 nm was assayed. The application thereof enabled the quantification of the protease of the present invention in a variety of specimens.

Example 14

(Purification of the protease using an antibody)

[0108] The obtained antibody was bound to a suitable immobilization carrier to prepare an affinity column, and the resulting column was used to purify the protease of the present invention. The affinity column was prepared by immobilizing an antibody using Cellulofine for NHS activation (Chisso Corporation) in accordance with the included instructions. The thus prepared swollen carrier (about 1 ml) was used to apply the culture supernatant in which the recombinant gene had been expressed in the 293 cell of the protease as described in Example 9. Thereafter, the column was washed with 50 mM Tris-HCl and 0.1M NaCl (pH 7.5, hereafter referred to as "TBS"), and elution was carried out using a urea-containing 0.1M glycine buffer (pH 3). The eluted fraction was neutralized with 1M Tris-HCl (pH 8.5) and then dialyzed against TBS. Fig. 20 shows the results of SDS-PAGE analysis of the resulting purified protease. Also, the resulting purified fraction was found to have vWF-cleaving activity. The cleavage point of the vWF fragmented by this recombinant protease was found to be the position between residues Tyr 842 and Met 843 based on the analysis of the N-terminal amino acid sequence of the fragment. Also established were clones (e.g., Clone Nos. CPHSWH-7.2 and 10) that could be similarly subjected to purification with the use of the monoclonal antibody prepared by the method as described in Example 13.

[0109] Subsequently, the partial amino acid sequence of the purified protease was determined. In accordance with a conventional technique, the protease was subjected to SDS-PAGE, transferred to a PVDF membrane, air-dried, and then subjected to analysis using an automated protein sequencer (model 492; PE Applied Biosystems). As a result, the protease was found to comprise Ala-Ala-Gly-Gly-Ile- as a partial N-terminal sequence. This sequence was congruous with the N-terminal sequence of the mature unit of the protease of the present invention, that was deduced from the genetic construction.

Example 15

(Neutralization of the protease activity using an antibody)

[0110] Activity of the aforementioned rabbit polyclonal antibody to neutralize the vWF-cleaving protease was evaluated. Normal rabbit serum, rabbit antiserum comprising the C-terminal peptide sequence (SEQ ID NO: 37), Phe-Ser-Pro-Ala-Pro-Gln-Pro-Arg-Arg-Leu-Leu-Pro-Gly-Pro-Gln-Glu-Asn-Ser-Val-Gln-Ser-Ser bound to KLH as an immunogen, and antiserum, the immunity of which had been induced by the protein expressed by the expression vector as shown in Example 7 or 8, were respectively allowed to pre-react at 37°C for 1 hour with 1 to 10 µg/ml of gene recombinant vWF-cleaving protease (approximated by the Bradford technique) at a volume ratio of 1:1. Alternatively, a 5-fold diluted antiserum was allowed to pre-react under the above conditions with the protease at a volume ratio of 1:1. Thereafter, vWF-cleaving activity was evaluated by the method described above. As a result, it was found that antiserum, which had activity of inhibiting the protease of the present invention, were prepared by immunizing the protein (Fig.21). (antagonist activity) (a metalloprotease inhibitor, i.e., EDTA, was determined to be a control). This indicates the possibility of constructing an acquired TTP patient-like model having a positive autoantibody against vWF-cleaving protease as well as the simple possibility of producing a neutralizing antibody.

Example 16

(Construction of C-terminus deleted modification unit)

[0111] Based on the strategy shown in Fig. 22, the full-length vWF-cleaving protease gene cloning vector (pCR 2.1 vWFCP) obtained in Example 6 or 7 was used to add a variant lacking domains located in a position following the C-terminus (T1135stop, W1016stop, W897stop, T581stop, and Q449stop: each numerical value indicates the number of amino acid residues between Met encoded by the initiation codon AGT and the termination codon, and indicates a site comprising the FLAG epitope (DNA sequence: gactacaaggacgatgacgataagtga (SEQ ID NO: 47) and amino acid sequence: Asp Tyr Lys Asp Asp Asp Lys (SEQ ID NO: 48)). Primers used herein are as follows. "S" indicates a sense primer, and "AS" indicates an antisense primer. Genes Stu I-S (SEQ ID NO: 38), Acc I-S (SEQ ID NO: 39), Avr II-S (SEQ ID NO: 40), Q449stop-AS (SEQ ID NO: 41), T581stop-AS (SEQ ID NO: 42), W897stop-AS (SEQ ID NO: 43), W1016stop-AS (SEQ ID NO: 44), T1135stop-AS (SEQ ID NO: 45), and full-length-AS (SEQ ID NO: 46) were prepared and incorporated in the pCAG expression vector in accordance with the method as used in Examples 8 and

9. This expression vector was introduced in the Hela cell. The primer pair shown at the bottom of the restriction map in the upper portion of Fig. 22 was used to obtain PCR fragments (A) to (F). Each PCR fragment was ligated to pCR 2.1 vWFCP. Further, the resultant was digested with *Stu*I/*Sal*I, and fragments (A) and (B) were digested with *Stu*I/*Sal*I and then ligated. These fragments were further digested with *Acc*I, and fragment (C) was also digested with *Acc*I, followed by ligation. The ligation product was digested with *Avr*II/*Sal*I, and fragments (D), (E), and (F) were also digested with *Avr*II/*Sal*I, followed by ligation. As a result, a variant lacking a region between the C-terminus and the position W897 was found to have activity, although it was the result of qualitative analysis. Such a way of approach enables the identification of various functional domains. The design of molecules comprising these domains and having no protease activity is considered to realize the design of antagonists or agonists.

Industrial Applicability

[0112] The findings of the present invention have led to the possibility of replacement therapy for patients having diseases resulting from deficiency of a protease, such as thrombotic thrombocytopenic purpura. This also realizes the establishment of methods for gene cloning and efficient purification from serum or plasma. In particular, the information provided by the present invention enables gene recombination based on the obtained nucleotide sequence and stable production and provision of the protease according to the present invention, which have been heretofore difficult to achieve. Also, these can be applied to replacement therapy for TTP patients, inhibition of platelet plug formation involved with heart infarction or brain infarction, inhibition of arteriosclerosis, prevention of restenosis, reembo-
lization, or infarction involved with PTCA, prevention of reembo-
lization involved with PTCR, and prevention of platelet plug formation caused by HUS or O-157. Diagnosis and therapy utilizing the gene encoding the protease of the present invention or an antibody thereagainst can be realized.

[0113] All publications cited herein are incorporated herein in their entirety. A person skilled in the art would easily understand that various modifications and changes of the present invention are feasible within the technical idea and the scope of the invention as disclosed in the attached claims. The present invention is intended to include such modifications and changes.

SEQUENCE LISTING

5 <110>JURIDICAL FOUNDATION THE CHEMO-SERO-THERAPEUTIC RESEARCH INSTITUTE
 <120>vWF-cleaving protease
 <130> PH1553-PCT
 10 <160>48

15 <210>1
 <211>5
 20 <212>PRT
 <213> Homo sapiens
 <400>1
 25 Leu Leu Val Ala Val
 1 5

30 <210>2
 <211>14
 35 <212>PRT
 <213> Homo sapiens
 <400>2
 40 Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val
 1 5 10

45 <210>3
 <211>161
 50 <212>PRT
 <213> Homo sapiens
 55 <400>3

	Ala	Ala	Gly	Gly	Ile	Leu	His	Leu	Glu	Leu	Leu	Val	Ala	Val	Gly
5	1				5					10					15
	Pro	Asp	Val	Phe	Gln	Ala	His	Gln	Lys	Asp	Thr	Glu	Arg	Tyr	Val
					20					25					30
10	Leu	Thr	Asn	Leu	Asn	Ile	Gly	Ala	Glu	Leu	Leu	Arg	Asp	Pro	Ser
					35					40					45
15	Leu	Gly	Ala	Gln	Phe	Arg	Val	His	Leu	Val	Lys	Met	Val	Ile	Leu
					50					55					60
20	Thr	Glu	Pro	Glu	Gly	Ala	Pro	Asn	Ile	Thr	Ala	Asn	Leu	Thr	Ser
					65					70					75
	Ser	Leu	Leu	Ser	Val	Cys	Gly	Trp	Ser	Gln	Thr	Ile	Asn	Pro	Glu
25					80					85					90
	Asp	Asp	Thr	Asp	Pro	Gly	His	Ala	Asp	Leu	Val	Leu	Tyr	Ile	Thr
					95					100					105
30	Arg	Phe	Asp	Leu	Glu	Leu	Pro	Asp	Gly	Asn	Arg	Gln	Val	Arg	Gly
					110					115					120
35	Val	Thr	Gln	Leu	Gly	Gly	Ala	Cys	Ser	Pro	Thr	Trp	Ser	Cys	Leu
					125					130					135
40	Ile	Thr	Glu	Asp	Thr	Gly	Phe	Asp	Leu	Gly	Val	Thr	Ile	Ala	His
					140					145					150
	Glu	Ile	Gly	His	Ser	Phe	Gly	Leu	Glu	His	Asp				
45					155					160					

<210>4

<211>15

<212>DNA

<213> Homo sapiens

<400>4

5 ctgctggtgg ccgtg

15

<210>5

10 <211>42

<212>DNA

15 <213> Homo sapiens

<400>5

20 gctgcaggcg gcatcctaca cctggagctg ctggaggccg tg

42

<210>6

25 <211>483

<212>DNA

30 <213> Homo sapiens.

<400>6

35 gctgcaggcg gcatcctaca cctggagctg ctggaggccg tgggccccga tgtcttccag 60

gctcaccaga aggacacaga gcgctatgtg ctaccaacc tcaacatcgg ggcagaactg 120

cttcgggacc cgtccctggg ggctcagtt cgggtgcacc tgggaagat ggtcattctg 180

40 acagagcctg aggggtgctc aaatatcaca gaaacctca cctcgtccct gctgagcgtc 240

tgtgggtgga gccagacat caacctgag gacgacacgg atcctggcca tgcagacctg 300

45 gtcctctata tcaactaggtt tgacctggag ttgcctgatg glaaccggca ggtgcggggc 360

gtcaccacgc tgggcgggtgc ctgctcccca acctggagct gcctcattac cgaggacact 420

ggcttcgacc tgggagtcac catctgcccac gagattgggc acagcttcgg cctggagcac 480

50 gac 483

<210>7

55 <211>161

<212>PRT

<213> Homo sapiens

 $\langle 400 \rangle_7$

gct gca ggc ggc atc cta cac ctg gag ctg ctg gtg gcc gtg ggc	45
Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly	
1 5 10 15	
ccc gat gtc ttc cag gct cac cag aag gac aca gag cgc tat gtg	90
Pro Asp Val Phe Gln Ala His Gln Lys Asp Thr Glu Arg Tyr Val	
20 25 30	
ctc acc aac ctc aac atc ggg gca gaa ctg ctt cgg gac ccg tcc	135
Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro Ser	
35 40 45	
ctg ggg gct cag ttt cgg gtg cac ctg gtg aag atg gtc att ctg	180
Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile Leu	
50 55 60	
aca gag cct gag ggt gct cca aat atc aca gca aac ctc acc tcg	225
Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr Ser	
65 70 75	
tcc ctg ctg agc gtc tgt ggg tgg agc cag acc atc aac cct gag	270
Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro Glu	
80 85 90	
gac gac acg gat cct ggc cat gct gac ctg gtc ctc tat atc act	315
Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile Thr	
95 100 105	
agg ttt gac ctg gag ttg cct gat ggt aac cgg cag gtg cgg ggc	360
Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg Gly	
110 115 120	
gtc acc cag ctg ggc ggt gcc tgc tcc cca acc tgg agc tgc ctc	405

Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys Leu
 5 125 130 135
 att acc gag gac act ggc ttc gac ctg gga gtc acc att gcc cat 450
 Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala His
 10 140 145 150
 gag att ggg cac agc ttc ggc ctg gag cac gac 483
 15 Glu Ile Gly His Ser Phe Gly Leu Glu His Asp
 155 160
 20 <210>8
 <211>29
 <212>PRT
 25 <213> Homo sapiens
 <400>8
 30 Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly
 1 5 10 15
 35 Pro Asp Val Phe Gln Ala His Gln Lys Asp Thr Arg Arg Tyr
 20 25
 40 <210>9
 <211>30
 <212>DNA
 45 <213> Homo sapiens
 <400>9
 50 gctgcaggcg gcatcctaca cctggagctg 30
 <210>10
 55 <211>21

<212>DNA

<213> Homo sapiens

<400>10

cccaatctca tgggcaatgg t

21

<210>11

<211>21

<212>DNA

<213> Homo sapiens

<400>11

cccaatctca tgggcaatgg t

21

<210>12

<211>30

<212>DNA

<213> Homo sapiens

<400>12

ccgatgttga ggttggtgag cacatagcgc

30

<210>13

<211>20

<212>DNA

<213> Homo sapiens

<400>13

gtgtcgtcct cagggttgat

20

<210>14

<211>21

<212>DNA

<213> Homo sapiens

<400>14

accatigccc atgagattgg g

21

<210>15

<211>4950

<212>DNA

<213> Homo sapiens

<400>15

aaccacgatg	tccttggcac	agcctctcat	ctgtcagatg	ggagcgggga	ccccggagag	60
ggagtcagcc	gaggctctgg	cattccctgt	gaacccccgt	ctgtgggttt	ctggctccagt	120
gtcccttctc	cagattagat	ggcttaggcc	tcctctaagg	gggtgggcgt	gcacatccgg	180
agagctgtct	gggtgacagg	actgggcctg	aggttaccct	gaactgcaac	catcttagag	240
caaggcccag	cttgcagcag	gaggagctgc	aggccgcccc	ccctagccac	ggccccctgcc	300
ctggcaggaa	gcttccaaga	gtaaacactg	cctaattcgt	ccgcccagta	gtgagcaggc	360
ctgtcccatt	ccatactgac	cagattccca	gtcaccaagg	ccccctctca	ctccgtctca	420
ctcctcgggc	tggctctcct	gaggatgcac	cagcgtcacc	cccgggcaag	atgccctccc	480
ctctgtgtgg	ccggaatcct	tgcctgtggc	tttctcctgg	gctgctgggg	accctcccat	540
ttccagcaga	gttgtcttca	ggctttggag	ccacaggccg	tgctctctta	cttgagccct	600
ggtgtctcct	taaaaggccg	ccctccttcc	cctggcttcc	agaggcagag	gcagaggcag	660
aggcgggctg	caggcggcat	cctacacctg	gagctgtctg	tggccgtggg	ccccgatgtc	720
ttccaggctc	accaggagga	cacagagcgc	tatgtgtctc	ccaacctcaa	catcggggca	780
gaactgcttc	gggacccgtc	cctgggggct	cagtttcggg	tgcacctggt	gaagatggtc	840
attctgacag	agcctgaggg	tgtctccaat	atcacagcca	acctcaccct	gtccctgctg	900
agcgtctgtg	ggaggagcca	gacctcaaac	cctgaggacg	acacggatcc	tggccatgct	960
gacctggtcc	tctatatac	taggtttgac	ctggagttgc	ctgatggtaa	ccggcaggtg	1020
cggggcgtca	cccagctggg	cgggtgcctgc	tccccaacct	ggagctgcct	cattaccgag	1080

gacac1ggct tgcacc1ggg agtcaccatt gcccatgaga t1gggcacag ct1cggcctg 1140
 5 gaggacgacg gcgcgcccgg cagcggctgc ggccccagcg gacacgtgat ggcttcggac 1200
 ggcgccgcgc cccgcgcccgg cctcgcc1gg tccccctgca gcccgccggca gctgctgagc 1260
 ctgctcagcg caggacgggc gcgctgcg1g tgggacccgc cgcggcc1ca acccgggtcc 1320
 10 gcggggcacc cgccggatgc gcagcc1ggc ctctactaca gcgccaacga gcag1gccgc 1380
 gtggcct1cg gcccgaaggc tgtcgcc1gc accttcgcca gggagcacct ggata1gtgc 1440
 15 caggccctct cctgccacac agaccgc1g gaccaaagca gctgcagccg cctcctcg1t 1500
 cctctcct1gg atgggacaga atgtggcg1g gagaagtgg1 gctccaaggg tgc1gcccgc 1560
 tccttgg1gg agctgacccc catagcagca gtgcatgggc gctgg1ctag ctgggg1ccc 1620
 20 cgaagtcc1t gctcccgc1c ctgcggagga ggtgtgg1ca ccaggaggcg gcag1gcaac 1680
 aaccccagac ctgcc1ttgg ggggcg1gca tgtgt1gg1g ctgacctca ggccgagatg 1740
 25 tgcaacac1c aggcctgcga gaagaccag ctggagt1ca tgtcgcaaca gtgcgccagg 1800
 accgacggcc agccgctgcg ctcc1ccct ggcggcgcct ccttctacca ctgggg1gct 1860
 gctgtaccac acagccaagg ggatgctctg tgcagacaca tgtgccgggc cattggcgag 1920
 30 agcttcatca tgaagcgtgg agacagct1c ctgatggga cccggtgtat gccaa1tggc 1980
 ccccgggagg accggaccct gagcctgtgt gt1tcgggca gctgcaggac atttggctgt 2040
 gatggtagga tggactccca gcaggtatgg gacagg1gcc aggtgtgtgg tggggacaac 2100
 35 agcacgtgca gccacggaa gggctct1tc acagctggca gagcgagaga atatgtcacg 2160
 tt1ctgacag ttacccccaa cctgaccagt gtctacattg ccaaccacag gcctctct1c 2220
 40 acacact1gg cgg1gaggat cggagggcgc tatgtcgtgg ctgggaagat gagcatctcc 2280
 cctaacacca cctaccc1c cctcctggag gatgg1cgtg tgcagtacag agtggccc1c 2340
 accgaggacc ggctgccccg cctggaggag atccgcatct ggggacccct ccaggaagat 2400
 45 gctgacatcc aggt1tacag gcggtatggc gaggagt1g gcaacc1cac ccgcccagac 2460
 atcacct1ca cctact1cca gcctaagcca cggcaggcct ggg1gtgggc cgctgtgcgt 2520
 50 gggccctgct cgg1gagctg tggggcaggg ctgcgctggg taaactacag ctgcctggac 2580
 caggccagga aggagt1gg1 ggagact1c cagtccaaag ggagccagca gccaccagcg 2640
 tggccagagg cctgcg1gct cgaaccc1gc cctccctact gggcggtggg agact1cggc 2700
 55 ccatgcagcg cctcct1tgg gggcgcc1g cgggagcggc cagtgcgctg cgtggaggcc 2760

	cagggcagcc	tcctgaagac	attgccccca	gcccggctgca	gagcaggggc	ccagcagcca	2820
5	gctgtggcgc	tggaaacctg	caacccccag	ccctgccctg	ccaggaggga	gggtgcagag	2880
	cccagctcat	gcacatcagc	tggtagagca	ggcctggcct	tggagaacga	gacctgtgtg	2940
	ccaggggcag	atggcctgga	ggctccagtg	acagaggggc	ctggctccgt	agatgagaag	3000
10	ctgcctgccc	ctgagccctg	tgtcgggag	tcatgtcctc	caggctgggg	ccatctggat	3060
	gccacctctg	caggggagaa	ggctccctcc	ccatggggca	gcatcaggac	gggggctcaa	3120
15	gctgcacacg	tgtggacccc	tgcggcaggg	tctgtctccg	tctcctgcgg	gcgaggtctg	3180
	atggagctgc	gtttcctgtg	catggactct	gcccicaggg	tgccgttcca	ggaagagctg	3240
	tgtggcctgg	caagcaagcc	tgggagccgg	cgggagggtc	gccaggctgt	cccgtgccct	3300
20	gctcgggtgg	agtacaagct	ggcggcctgc	agcgtgagct	gtgggagagg	ggctgtcggg	3360
	aggatcctgt	attgtgcccc	ggcccatggg	gaggacgatg	gtgaggagat	cctgttggac	3420
25	acccagtgcc	aggggctgcc	tgcgccgaa	ccccaggagg	cctgcagcct	ggagccctgc	3480
	ccacctaggt	ggaaagtcat	gtcccttggc	ccatgttcgg	ccagctgtgg	ccttggcact	3540
	gctagacgct	cggtagcctg	tgtgcagctc	gaccaaggcc	aggacgtgga	ggtaggacgag	3600
30	gcggcctgtg	cggcgctggt	gcggcccag	gccagtgtcc	cctgtctcat	tgccgactgc	3660
	acctaccgct	ggcatgttgg	cacctggatg	gagtgtcttg	tttcctgtgg	ggatggcatc	3720
35	cagcgccggc	gtgacacctg	cctcggaccc	caggcccagg	cgcctgtgcc	agctgatctc	3780
	tgccagcact	tgcccaagcc	ggtgactgtg	cgtggctgct	gggctgggcc	ctgtgtggga	3840
	cagggtagcg	ccagcctggg	gccccacgaa	gaagccgctg	ctccaggacg	gaccacagcc	3900
40	acccctgctg	gtgcctccct	ggagtggctc	caggcccggg	gcctgtctct	ctccccggct	3960
	ccccagcctc	ggcggctcct	gcccggggcc	caggaaaact	cagtgcagtc	cagtgcctgt	4020
	ggcaggcagc	accttgagcc	aacaggaacc	attgacatgc	gaggcccagg	gcaggcagac	4080
45	tgtgcagtgg	ccattgggcg	gcccctcggg	gaggtagtga	ccctccgcgt	ccttgagagt	4140
	tctctcaact	gcagtgcggg	ggacatgttg	ctgctttggg	gccggctcac	ctggaggaag	4200
50	atgtgcagga	agctgttggg	catgactttc	agctccaaga	ccaacacgct	ggtaggtgagg	4260
	cagcgctgcg	ggcggccagg	aggtggggtg	ctgctgcggg	atgggagcca	gcttgcctct	4320
	gaaaccttct	acagagaatg	tgacatgcag	ctctttgggc	cctggggtag	aatcgtgagc	4380
55	ccctcgttga	gtccagccac	gagtaatgca	gggggctgcc	ggctcttcat	taatgtggct	4440

5 cgcacgcac ggattgccat ccattgccctg gccaccaaca tgggcgcctgg gaccgagggg 4500
 gccaatgcc a gctacatcct gatccgggac acccacagct tgaggaccac agcgttccat 4560
 gggcagcagg tgctctactg ggagtcagag agcagccagg ctgagatgga gttcagcgag 4620
 10 ggcttccatga aggttcaggc cagcctgcgg ggccagtact ggacctcca atcatgggta 4680
 ccggagatgc aggacctca gtccctggaag ggaaaggaag gaacctgagg gtcattgaac 4740
 atttggtccg tgcctggcca gccctggagg gttgacctt ggtctcagt ctttccaatt 4800
 15 cgaacttttt ccaatcttag gtatctactt tagagctctt tccaatgtcc aaaaggctag 4860
 ggggttgag gtggggactc tggaaaagca gccccattt cctcgggtac caataaataa 4920
 aacatgcagg ccaaaaaaaaa aaaaaaaaaa 4950

<210>16

<211>1353

<212>PRT

<213> Homo sapiens

<400>16

30 gct gca ggc ggc atc cta cac ctg gag ctg ctg gtg gcc gtg ggc 45
 Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly
 35 1 5 10 15
 ccc gat gtc ttc cag gct cac cag gag gac aca gag cgc tat gtg 90
 40 Pro Asp Val Phe Gln Ala His Gln Glu Asp Thr Glu Arg Tyr Val
 20 25 30
 ctc acc aac ctc aac atc ggg gca gaa ctg ctt cgg gac ccg tcc 135
 45 Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro Ser
 35 40 45
 ctg ggg gct cag ttt cgg gtg cac ctg gtg aag atg gtc att ctg 180
 50 Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile Leu
 50 55 60
 55 aca gag cct gag ggt gct cca aat atc aca gcc aac ctc acc tcg 225

	Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr Ser	
5	65 70 75	
	tcc ctg ctg agc gtc tgt ggg tgg agc cag acc atc aac cct gag	270
10	Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro Glu	
	80 85 90	
	gac gac acg gat cct ggc cat gct gac ctg gtc ctc tat atc act	315
15	Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile Thr	
	95 100 105	
20	agg ttt gac ctg gag ttg cct gat ggt aac cgg cag gtg cgg ggc	360
	Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg Gly	
	110 115 120	
25	gtc acc cag ctg ggc ggt gcc tgc tcc cca acc tgg agc tgc ctc	405
	Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys Leu	
	125 130 135	
30	att acc gag gac act ggc ttc gac ctg gga gtc acc att gcc cat	450
	Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala His	
	140 145 150	
35	gag att ggg cac agc ttc ggc ctg gag cac gac ggc gcg ccc ggc	495
	Glu Ile Gly His Ser Phe Gly Leu Glu His Asp Gly Ala Pro Gly	
40	155 160 165	
	agc ggc tgc ggc ccc agc gga cac gtg atg gct tgc gac ggc gcc	540
	Ser Gly Cys Gly Pro Ser Gly His Val Met Ala Ser Asp Gly Ala	
45	170 175 180	
	gcg ccc cgc gcc ggc ctc gcc tgg tcc ccc tgc agc cgc cgg cag	585
50	Ala Pro Arg Ala Gly Leu Ala Trp Ser Pro Cys Ser Arg Arg Gln	
	185 190 195	
55	ctg ctg agc ctg ctc agc gca gga cgg gcg cgc tgc gtg tgg gac	630
	Leu Leu Ser Leu Leu Ser Ala Gly Arg Ala Arg Cys Val Trp Asp	

	200	205	210	
5	ccg ccg cgg cct caa ccc ggg tcc gcg ggg cac ccg ccg gat gcg			675
	Pro Pro Arg Pro Gln Pro Gly Ser Ala Gly His Pro Pro Asp Ala			
	215	220	225	
10	cag cct ggc ctc tac tac agc gcc aac gag cag tgc cgc gtg gcc			720
	Gln Pro Gly Leu Tyr Tyr Ser Ala Asn Glu Gln Cys Arg Val Ala			
15	230	235	240	
	ttc ggc ccc aag gct gtc gcc tgc acc ttc gcc agg gag cac ctg			765
	Phe Gly Pro Lys Ala Val Ala Cys Thr Phe Ala Arg Glu His Leu			
20	245	250	255	
	gat atg tgc cag gcc ctc tcc tgc cac aca gac ccg ctg gac caa			810
25	Asp Met Cys Gln Ala Leu Ser Cys His Thr Asp Pro Leu Asp Gln			
	260	265	270	
	agc agc tgc agc cgc ctc ctc gtt cct ctc ctg gat ggg aca gaa			855
30	Ser Ser Cys Ser Arg Leu Leu Val Pro Leu Leu Asp Gly Thr Glu			
	275	280	285	
35	tgt ggc gtg gag aag tgg tgc tcc aag ggt cgc tgc cgc tcc ctg			900
	Cys Gly Val Glu Lys Trp Cys Ser Lys Gly Arg Cys Arg Ser Leu			
	290	295	300	
40	gtg gag ctg acc ccc ata gca gca gtg cat ggg cgc tgg tct agc			945
	Val Glu Leu Thr Pro Ile Ala Ala Val His Gly Arg Trp Ser Ser			
	305	310	315	
45	tgg ggt ccc cga agt cct tgc tcc cgc tcc tgc gga gga ggt gtg			990
	Trp Gly Pro Arg Ser Pro Cys Ser Arg Ser Cys Gly Gly Gly Val			
	320	325	330	
50	gtc acc agg agg cgg cag tgc aac aac ccc aga cct gcc ttt ggg			1035
	Val Thr Arg Arg Arg Gln Cys Asn Asn Pro Arg Pro Ala Phe Gly			
55	335	340	345	

	ggg cgt gca tgt gtt ggt gct gac ctc cag gcc gag atg tgc aac	1080
5	Gly Arg Ala Cys Val Gly Ala Asp Leu Gln Ala Glu Met Cys Asn	
	350 355 360	
	act cag gcc tgc gag aag acc cag ctg gag ttc atg tcg caa cag	1125
10	Thr Gln Ala Cys Glu Lys Thr Gln Leu Glu Phe Met Ser Gln Gln	
	365 370 375	
	tgc gcc agg acc gac ggc cag ccg ctg cgc tcc tcc cct ggc ggc	1170
15	Cys Ala Arg Thr Asp Gly Gln Pro Leu Arg Ser Ser Pro Gly Gly	
	380 385 390	
	gcc tcc ttc tac cac tgg ggt gct gct gta cca cac agc caa ggg	1215
20	Ala Ser Phe Tyr His Trp Gly Ala Ala Val Pro His Ser Gln Gly	
	395 400 405	
25	gat gct ctg tgc aga cac atg tgc cgg gcc att ggc gag agc ttc	1260
	Asp Ala Leu Cys Arg His Met Cys Arg Ala Ile Gly Glu Ser Phe	
30	410 415 420	
	atc atg aag cgt gga gac agc ttc ctc gat ggg acc cgg tgt atg	1305
	Ile Met Lys Arg Gly Asp Ser Phe Leu Asp Gly Thr Arg Cys Met	
35	425 430 435	
	cca agt ggc ccc cgg gag gac ggg acc ctg agc ctg tgt gtg tcg	1350
40	Pro Ser Gly Pro Arg Glu Asp Gly Thr Leu Ser Leu Cys Val Ser	
	440 445 450	
	ggc agc tgc agg aca ttt ggc tgt gat ggt agg atg gac tcc cag	1395
45	Gly Ser Cys Arg Thr Phe Gly Cys Asp Gly Arg Met Asp Ser Gln	
	455 460 465	
	cag gta tgg gac agg tgc cag gtg tgt ggt ggg gac aac agc acg	1440
50	Gln Val Trp Asp Arg Cys Gln Val Cys Gly Gly Asp Asn Ser Thr	
	470 475 480	
55	tgc agc cca cgg aag ggc tct ttc aca gct ggc aga gcg aga gaa	1485

	Cys Ser Pro Arg Lys Gly Ser Phe Thr Ala Gly Arg Ala Arg Glu	
5	485 490 495	
	tat gtc acg ttt ctg aca gtt acc ccc aac ctg acc agt gtc tac	1530
	Tyr Val Thr Phe Leu Thr Val Thr Pro Asn Leu Thr Ser Val Tyr	
10	500 505 510	
	att gcc aac cac agg cct ctc ttc aca cac ttg gcg gtg agg atc	1575
15	Ile Ala Asn His Arg Pro Leu Phe Thr His Leu Ala Val Arg Ile	
	515 520 525	
	gga ggg cgc tat gtc gtg gct ggg aag atg agc atc tcc cct aac	1620
20	Gly Gly Arg Tyr Val Val Ala Gly Lys Met Ser Ile Ser Pro Asn	
	530 535 540	
	acc acc tac ccc tcc ctc ctg gag gat ggt cgt gtc gag tac aga	1665
25	Thr Thr Tyr Pro Ser Leu Leu Glu Asp Gly Arg Val Glu Tyr Arg	
	545 550 555	
30	gtg gcc ctc acc gag gac cgg ctg ccc cgc ctg gag gag atc cgc	1710
	Val Ala Leu Thr Glu Asp Arg Leu Pro Arg Leu Glu Glu Ile Arg	
	560 565 570	
35	atc tgg gga ccc ctc cag gaa gat gct gac atc cag gtt tac agg	1755
	Ile Trp Gly Pro Leu Gln Glu Asp Ala Asp Ile Gln Val Tyr Arg	
40	575 580 585	
	cgg tat ggc gag gag tat ggc aac ctc acc cgc cca gac atc acc	1800
	Arg Tyr Gly Glu Glu Tyr Gly Asn Leu Thr Arg Pro Asp Ile Thr	
45	590 595 600	
	ttc acc tac ttc cag cct aag cca cgg cag gcc tgg gtg tgg gcc	1845
50	Phe Thr Tyr Phe Gln Pro Lys Pro Arg Gln Ala Trp Val Trp Ala	
	605 610 615	
	gct gtg cgt ggg ccc tgc tcg gtg agc lgt ggg gca ggg ctg cgc	1890
55	Ala Val Arg Gly Pro Cys Ser Val Ser Cys Gly Ala Gly Leu Arg	

	620	625	630	
5	tgg gta aac tac agc tgc ctg gac cag gcc agg aag gag ttg gtg			1935
	Trp Val Asn Tyr Ser Cys Leu Asp Gln Ala Arg Lys Glu Leu Val			
	635	640	645	
10	gag act gtc cag tgc caa ggg agc cag cag cca cca gcg tgg cca			1980
	Glu Thr Val Gln Cys Gln Gly Ser Gln Gln Pro Pro Ala Trp Pro			
	650	655	660	
15	gag gcc tgc gtg ctc gaa ccc tgc cct ccc tac tgg gcg gtg gga			2025
	Glu Ala Cys Val Leu Glu Pro Cys Pro Pro Tyr Trp Ala Val Gly			
	665	670	675	
20	gac ttc ggc cca tgc agc gcc tcc tgt ggg ggc ggc ctg cgg gag			2070
	Asp Phe Gly Pro Cys Ser Ala Ser Cys Gly Gly Gly Leu Arg Glu			
25	680	685	690	
	cgg cca gtg cgc tgc gtg gag gcc cag ggc agc ctc ctg aag aca			2115
30	Arg Pro Val Arg Cys Val Glu Ala Gln Gly Ser Leu Leu Lys Thr			
	695	700	705	
	ttg ccc cca gcc cgg tgc aga gca ggg gcc cag cag cca gct gtg			2160
35	Leu Pro Pro Ala Arg Cys Arg Ala Gly Ala Gln Gln Pro Ala Val			
	710	715	720	
40	gcg ctg gaa acc tgc aac ccc cag ccc tgc cct gcc agg tgg gag			2205
	Ala Leu Glu Thr Cys Asn Pro Gln Pro Cys Pro Ala Arg Trp Glu			
	725	730	735	
45	gtg tca gag ccc agc tca tgc aca tca gct ggt gga gca ggc ctg			2250
	Val Ser Glu Pro Ser Ser Cys Thr Ser Ala Gly Gly Ala Gly Leu			
	740	745	750	
50	gcc ttg gag aac gag acc tgt gtg cca ggg gca gat ggc ctg gag			2295
	Ala Leu Glu Asn Glu Thr Cys Val Pro Gly Ala Asp Gly Leu Glu			
55	755	760	765	

	gct cca gtg act gag ggg cct ggc tcc gta gat gag aag ctg cct	2340
5	Ala Pro Val Thr Glu Gly Pro Gly Ser Val Asp Glu Lys Leu Pro	
	770 775 780	
	gcc cct gag ccc tgt gtc ggg atg tca tgt cct cca ggc tgg ggc	2385
10	Ala Pro Glu Pro Cys Val Gly Met Ser Cys Pro Pro Gly Trp Gly	
	785 790 795	
	cat ctg gat gcc acc tct gca ggg gag aag gct ccc tcc cca tgg	2430
15	His Leu Asp Ala Thr Ser Ala Gly Glu Lys Ala Pro Ser Pro Trp	
	800 805 810	
	ggc agc atc agg acg ggg gct caa gct gca cac gtg tgg acc cct	2475
20	Gly Ser Ile Arg Thr Gly Ala Gln Ala Ala His Val Trp Thr Pro	
	815 820 825	
	gcg gca ggg tgg tgc tcc gtc tcc tgc ggg cga ggt ctg atg gag	2520
25	Ala Ala Gly Ser Cys Ser Val Ser Cys Gly Arg Gly Leu Met Glu	
	830 835 840	
	ctg cgt ttc ctg tgc atg gac tct gcc ctg agg gtg cct gtc cag	2565
30	Leu Arg Phe Leu Cys Met Asp Ser Ala Leu Arg Val Pro Val Gln	
	845 850 855	
	gaa gag ctg tgt ggc ctg gca agc aag cct ggg agc cgg cgg gag	2610
35	Glu Glu Leu Cys Gly Leu Ala Ser Lys Pro Gly Ser Arg Arg Glu	
	860 865 870	
	gtc tgc cag gct gtc ccg tgc cct gct cgg tgg cag tac aag ctg	2655
40	Val Cys Gln Ala Val Pro Cys Pro Ala Arg Trp Gln Tyr Lys Leu	
	875 880 885	
	gcg gcc tgc agc gtg agc tgt ggg aga ggg gtc gtg cgg agg atc	2700
45	Ala Ala Cys Ser Val Ser Cys Gly Arg Gly Val Val Arg Arg Ile	
	890 895 900	
	ctg tat tgt gcc cgg gcc cat ggg gag gac gat ggt gag gag atc	2745
50		
55		

	Leu Tyr Cys Ala Arg Ala His Gly Glu Asp Asp Gly Glu Glu Ile	
5	905 910 915	
	ctg ttg gac acc cag tgc cag ggg ctg cct cgc ccg gaa ccc cag	2790
10	Leu Leu Asp Thr Gln Cys Gln Gly Leu Pro Arg Pro Glu Pro Gln	
	920 925 930	
	gag gcc tgc agc ctg gag ccc tgc cca cct agg tgg aaa gtc atg	2835
15	Glu Ala Cys Ser Leu Glu Pro Cys Pro Pro Arg Trp Lys Val Met	
	935 940 945	
20	tcc ctt ggc cca tgt tgc gcc agc tgt ggc ctt ggc act gct aga	2880
	Ser Leu Gly Pro Cys Ser Ala Ser Cys Gly Leu Gly Thr Ala Arg	
	950 955 960	
25	cgc tgc gtg gcc tgt gtg cag ctc gac caa ggc cag gac gtg gag	2925
	Arg Ser Val Ala Cys Val Gln Leu Asp Gln Gly Gln Asp Val Glu	
	965 970 975	
30	gtg gac gag gcg gcc tgt gcg gcg ctg gtg cgg ccc gag gcc agt	2970
	Val Asp Glu Ala Ala Cys Ala Ala Leu Val Arg Pro Glu Ala Ser	
	980 985 990	
35	gtc ccc tgt ctc att gcc gac tgc acc tac cgc tgg cat gtt ggc	3015
	Val Pro Cys Leu Ile Ala Asp Cys Thr Tyr Arg Trp His Val Gly	
40	995 1000 1005	
	acc tgg atg gag tgc tct gtt tcc tgt ggg gat ggc atc cag cgc	3060
	Thr Trp Met Glu Cys Ser Val Ser Cys Gly Asp Gly Ile Gln Arg	
45	1010 1015 1020	
	cgg cgt gac acc tgc ctc gga ccc cag gcc cag gcg cct gtg cca	3105
50	Arg Arg Asp Thr Cys Leu Gly Pro Gln Ala Gln Ala Pro Val Pro	
	1025 1030 1035	
	gct gat ttc tgc cag cac ttg ccc aag ccg gtg act gtg cgt ggc	3150
55	Ala Asp Phe Cys Gln His Leu Pro Lys Pro Val Thr Val Arg Gly	

	1040	1045	1050	
5	tgc tgg gct ggg ccc tgt gtg gga cag ggt acg ccc agc ctg gtg			3195
	Cys Trp Ala Gly Pro Cys Val Gly Gln Gly Thr Pro Ser Leu Val			
	1055	1060	1065	
10	ccc cac gaa gaa gcc gct gct cca gga cgg acc aca gcc acc cct			3240
	Pro His Glu Glu Ala Ala Ala Pro Gly Arg Thr Thr Ala Thr Pro			
	1070	1075	1080	
15	gct ggt gcc tcc ctg gag tgg tcc cag gcc cgg ggc ctg etc ttc			3285
	Ala Gly Ala Ser Leu Glu Trp Ser Gln Ala Arg Gly Leu Leu Phe			
	1085	1090	1095	
20	tcc ccg gct ccc cag cct cgg cgg ctc ctg ccc ggg ccc cag gaa			3330
	Ser Pro Ala Pro Gln Pro Arg Arg Leu Leu Pro Gly Pro Gln Glu			
25	1100	1105	1110	
	aac tca gtg cag tcc agt gcc tgt ggc agg cag cac ctt gag cca			3375
30	Asn Ser Val Gln Ser Ser Ala Cys Gly Arg Gln His Leu Glu Pro			
	1115	1120	1125	
35	aca gga acc att gac atg cga ggc cca ggg cag gca gac tgt gca			3420
	Thr Gly Thr Ile Asp Met Arg Gly Pro Gly Gln Ala Asp Cys Ala			
	1130	1135	1140	
40	gtg gcc att ggg cgg ccc ctc ggg gag gtg gtg acc ctc cgc gtc			3465
	Val Ala Ile Gly Arg Pro Leu Gly Glu Val Val Thr Leu Arg Val			
	1145	1150	1155	
45	ctt gag agt tct ctc aac tgc agt gcg ggg gac atg ttg ctg ctt			3510
	Leu Glu Ser Ser Leu Asn Cys Ser Ala Gly Asp Met Leu Leu Leu			
	1160	1165	1170	
50	tgg ggc cgg ctc acc tgg agg aag atg tgc agg aag ctg ttg gac			3555
	Trp Gly Arg Leu Thr Trp Arg Lys Met Cys Arg Lys Leu Leu Asp			
55	1175	1180	1185	

	atg act ttc agc tcc aag acc aac acg ctg gtg gtg agg cag cgc	3600
5	Met Thr Phe Ser Ser Lys Thr Asn Thr Leu Val Val Arg Gln Arg	
	1190 1195 1200	
10	tgc ggg cgg cca gga ggt ggg gtg ctg ctg cgg tat ggg agc cag	3645
	Cys Gly Arg Pro Gly Gly Gly Val Leu Leu Arg Tyr Gly Ser Gln	
	1205 1210 1215	
15	ctt gct cct gaa acc ttc tac aga gaa tgt gac atg cag ctc ttt	3690
	Leu Ala Pro Glu Thr Phe Tyr Arg Glu Cys Asp Met Gln Leu Phe	
	1220 1225 1230	
20	ggg ccc tgg ggt gaa atc gtg agc ccc tcg ctg agt cca gcc acg	3735
	Gly Pro Trp Gly Glu Ile Val Ser Pro Ser Leu Ser Pro Ala Thr	
	1235 1240 1245	
25	agt aat gca ggg ggc tgc cgg ctc ttc att aat gtg gct ccg cac	3780
	Ser Asn Ala Gly Gly Cys Arg Leu Phe Ile Asn Val Ala Pro His	
30	1250 1255 1260	
	gca cgg att gcc atc cat gcc ctg gcc acc aac atg ggc gct ggg	3825
	Ala Arg Ile Ala Ile His Ala Leu Ala Thr Asn Met Gly Ala Gly	
35	1265 1270 1275	
	acc gag gga gcc aat gcc agc tac atc ttg atc cgg gac acc cac	3870
40	Thr Glu Gly Ala Asn Ala Ser Tyr Ile Leu Ile Arg Asp Thr His	
	1280 1285 1290	
45	agc ttg agg acc aca gcg ttc cat ggg cag cag gtg ctc tac tgg	3915
	Ser Leu Arg Thr Thr Ala Phe His Gly Gln Gln Val Leu Tyr Trp	
	1295 1300 1305	
50	gag tca gag agc agc cag gct gag atg gag ttc agc gag ggc ttc	3960
	Glu Ser Glu Ser Ser Gln Ala Glu Met Glu Phe Ser Glu Gly Phe	
	1310 1315 1320	
55	ctg aag gct cag gcc agc ctg cgg ggc cag tac tgg acc ctc caa	4005

Leu Lys Ala Gln Ala Ser Leu Arg Gly Gln Tyr Trp Thr Leu Gln
 5 1325 1330 1335
 tca tgg gla ccg gag atg cag gac cct cag tcc tgg aag gga aag 4050
 Ser Trp Val Pro Glu Met Gln Asp Pro Gln Ser Trp Lys Gly Lys
 10 1340 1345 1350
 gaa gga acc 4059
 Glu Gly Thr
 15
 <210>17
 20 <211>1297
 <212>PRT
 <213> Homo sapiens
 25 <400>17
 gct gca ggc ggc atc cta cac ctg gag ctg ctg gtg gcc gtg ggc 45
 30 Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly
 1 5 10 15
 ccc gat gtc ttc cag gct cac cag gag gac aca gag cgc tat gtg 90
 35 Pro Asp Val Phe Gln Ala His Gln Glu Asp Thr Glu Arg Tyr Val
 20 25 30
 40 ctc acc aac ctc aac atc ggg gca gaa ctg ctt cgg gac ccg tcc 135
 Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro Ser
 35 40 45
 45 ctg ggg gct cag ttt cgg gtg cac ctg gtg aag atg gtc att ctg 180
 Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile Leu
 50 50 55 60
 aca gag cct gag ggt gct cca aat atc aca gcc aac ctc acc tcg 225
 Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr Ser
 55 65 70 75

	tcc ctg ctg agc gtc tgt ggg tgg agc cag acc atc aac cct gag	270
5	Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro Glu	
	80 85 90	
	gac gac acg gat cct ggc cat gct gac ctg gtc ctc tat atc act	315
10	Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile Thr	
	95 100 105	
15	agg ttt gac ctg gag ttg cct gat ggt aac cgg cag gtg cgg ggc	360
	Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg Gly	
	110 115 120	
20	gtc acc cag ctg ggc ggt gcc tgc tcc cca acc tgg agc tgc ctc	405
	Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys Leu	
	125 130 135	
25	att acc gag gac act ggc ttc gac ctg gga gtc acc att gcc cat	450
	Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala His	
30	140 145 150	
	gag att ggg cac agc ttc ggc ctg gag cac gac ggc gcg ccc ggc	495
	Glu Ile Gly His Ser Phe Gly Leu Glu His Asp Gly Ala Pro Gly	
35	155 160 165	
	agc ggc tgc ggc ccc agc gga cac gtg atg gct tgc gac ggc gcc	540
40	Ser Gly Cys Gly Pro Ser Gly His Val Met Ala Ser Asp Gly Ala	
	170 175 180	
	gcg ccc cgc gcc ggc ctc gcc tgg tcc ccc tgc agc cgc cgg cag	585
45	Ala Pro Arg Ala Gly Leu Ala Trp Ser Pro Cys Ser Arg Arg Gln	
	185 190 195	
50	ctg ctg agc ctg ctc agc gca gga cgg gcg cgc tgc gtg tgg gac	630
	Leu Leu Ser Leu Leu Ser Ala Gly Arg Ala Arg Cys Val Trp Asp	
	200 205 210	
55	ccg ccg cgg cct caa ccc ggg tcc gcg ggg cac ccg ccg gat gcg	675

	Pro Pro Arg Pro Gln Pro Gly Ser Ala Gly His Pro Pro Asp Ala	
5	215 220 225	
	cag cct ggc ctc tac tac agc gcc aac gag cag tgc cgc gtg gcc	720
	Gln Pro Gly Leu Tyr Tyr Ser Ala Asn Glu Gln Cys Arg Val Ala	
10	230 235 240	
	ttc ggc ccc aag gct gtc gcc tgc acc ttc gcc agg gag cac ctg	765
	Phe Gly Pro Lys Ala Val Ala Cys Thr Phe Ala Arg Glu His Leu	
15	245 250 255	
	gat atg tgc cag gcc ctc tcc tgc cac aca gac ccg ctg gac caa	810
20	Asp Met Cys Gln Ala Leu Ser Cys His Thr Asp Pro Leu Asp Gln	
	260 265 270	
	agc agc tgc agc cgc ctc ctc gtt cct ctc ctg gat ggg aca gaa	855
25	Ser Ser Cys Ser Arg Leu Leu Val Pro Leu Leu Asp Gly Thr Glu	
	275 280 285	
	tgt ggc gtg gag aag tgg tgc tcc aag ggt cgc tgc cgc tcc ctg	900
30	Cys Gly Val Glu Lys Trp Cys Ser Lys Gly Arg Cys Arg Ser Leu	
	290 295 300	
	gtg gag ctg acc ccc ata gca gca gtg cat ggg cgc tgg tct agc	945
35	Val Glu Leu Thr Pro Ile Ala Ala Val His Gly Arg Trp Ser Ser	
	305 310 315	
	tgg ggt ccc cga agt cct tgc tcc cgc tcc tgc gga gga ggt gtg	990
40	Trp Gly Pro Arg Ser Pro Cys Ser Arg Ser Cys Gly Gly Gly Val	
	320 325 330	
	gtc acc agg agg cgg cag tgc aac aac ccc aga cct gcc ttt ggg	1035
50	Val Thr Arg Arg Arg Gln Cys Asn Asn Pro Arg Pro Ala Phe Gly	
	335 340 345	
	ggg cgt gca tgt gtt ggt gct gac ctc cag gcc gag atg tgc aac	1080
55	Gly Arg Ala Cys Val Gly Ala Asp Leu Gln Ala Glu Met Cys Asn	

	350	355	360	
5	act cag gcc tgc gag aag acc cag ctg gag ttc atg tcg caa cag			1125
	Thr Gln Ala Cys Glu Lys Thr Gln Leu Glu Phe Met Ser Gln Gln			
	365	370	375	
10	tgc gcc agg acc gac ggc cag ccg ctg cgc tcc tcc cct ggc ggc			1170
	Cys Ala Arg Thr Asp Gly Gln Pro Leu Arg Ser Ser Pro Gly Gly			
	380	385	390	
15	gcc tcc ttc tac cac tgg ggt gct gct gta cca cac agc caa ggg			1215
	Ala Ser Phe Tyr His Trp Gly Ala Ala Val Pro His Ser Gln Gly			
20	395	400	405	
	gat gct ctg tgc aga cac atg tgc cgg gcc att ggc gag agc ttc			1260
25	Asp Ala Leu Cys Arg His Met Cys Arg Ala Ile Gly Glu Ser Phe			
	410	415	420	
	atc atg aag cgt gga gac agc ttc ctg gat ggg acc cgg tgt atg			1305
30	Ile Met Lys Arg Gly Asp Ser Phe Leu Asp Gly Thr Arg Cys Met			
	425	430	435	
	cca agt ggc ccc cgg gag gac ggg acc ctg agc ctg tgt gtg tcg			1350
35	Pro Ser Gly Pro Arg Glu Asp Gly Thr Leu Ser Leu Cys Val Ser			
	440	445	450	
40	ggc agc tgc agg aca ttt ggc tgt gat ggt agg atg gac tcc cag			1395
	Gly Ser Cys Arg Thr Phe Gly Cys Asp Gly Arg Met Asp Ser Gln			
	455	460	465	
45	cag gla tgg gac agg tgc cag gtg tgt ggt ggg gac aac agc acg			1440
	Gln Val Trp Asp Arg Cys Gln Val Cys Gly Gly Asp Asn Ser Thr			
	470	475	480	
50	tgc agc cca cgg aag ggc tct ttc aca gct ggc aga gcg aga gaa			1485
	Cys Ser Pro Arg Lys Gly Ser Phe Thr Ala Gly Arg Ala Arg Glu			
55	485	490	495	

	tat gtc acg ttt ctg aca gtt acc ccc aac ctg acc agt gtc tac	1530
5	Tyr Val Thr Phe Leu Thr Val Thr Pro Asn Leu Thr Ser Val Tyr	
	500 505 510	
	att gcc aac cac agg cct ctc ttc aca cac ttg gcg gtg agg atc	1575
10	Ile Ala Asn His Arg Pro Leu Phe Thr His Leu Ala Val Arg Ile	
	515 520 525	
	gga ggg cgc tat gtc gtg gct ggg aag atg agc atc tcc cct aac	1620
15	Gly Gly Arg Tyr Val Val Ala Gly Lys Met Ser Ile Ser Pro Asn	
	530 535 540	
	acc acc tac ccc tcc ctc ctg gag gat ggt cgt gtc gag tac aga	1665
20	Thr Thr Tyr Pro Ser Leu Leu Glu Asp Gly Arg Val Glu Tyr Arg	
	545 550 555	
	gtg gcc ctc acc gag gac cgg ctg ccc cgc ctg gag gag atc cgc	1710
25	Val Ala Leu Thr Glu Asp Arg Leu Pro Arg Leu Glu Glu Ile Arg	
	560 565 570	
	atc tgg gga ccc ctc cag gaa gat gct gac atc cag gtt tac agg	1755
30	Ile Trp Gly Pro Leu Gln Glu Asp Ala Asp Ile Gln Val Tyr Arg	
	575 580 585	
	cgg tat ggc gag gag tat ggc aac ctc acc cgc cca gac atc acc	1800
35	Arg Tyr Gly Glu Glu Tyr Gly Asn Leu Thr Arg Pro Asp Ile Thr	
	590 595 600	
	ttc acc tac ttc cag cct aag cca cgg cag gcc tgg gtg tgg gcc	1845
40	Phe Thr Tyr Phe Gln Pro Lys Pro Arg Gln Ala Trp Val Trp Ala	
	605 610 615	
	gct gtg cgt ggg ccc tgc tgc gtg agc tgt ggg gca ggg ctg cgc	1890
45	Ala Val Arg Gly Pro Cys Ser Val Ser Cys Gly Ala Gly Leu Arg	
	620 625 630	
	tgg gta aac tac agc tgc ctg gac cag gcc agg aag gag ttg gtg	1935
50		
55		

	Trp Val Asn Tyr Ser Cys Leu Asp Gln Ala Arg Lys Glu Leu Val	
5	635 640 645	
	gag act gtc cag tgc caa ggg agc cag cag cca cca gcg tgg cca	1980
	Glu Thr Val Gln Cys Gln Gly Ser Gln Gln Pro Pro Ala Trp Pro	
10	650 655 660	
	gag gcc tgc gtg ctg gaa ccc tgc cct ccc tac tgg gcg gtg gga	2025
15	Glu Ala Cys Val Leu Glu Pro Cys Pro Pro Tyr Trp Ala Val Gly	
	665 670 675	
	gac ttc ggc cca tgc agc gcc tcc tgt ggg ggc ggc ctg cgg gag	2070
20	Asp Phe Gly Pro Cys Ser Ala Ser Cys Gly Gly Gly Leu Arg Glu	
	680 685 690	
25	cgg cca gtg cgc tgc gtg gag gcc cag ggc agc ctg aag aca	2115
	Arg Pro Val Arg Cys Val Glu Ala Gln Gly Ser Leu Leu Lys Thr	
	695 700 705	
30	ttg ccc cca gcc cgg tgc aga gca ggg gcc cag cag cca gct gtg	2160
	Leu Pro Pro Ala Arg Cys Arg Ala Gly Ala Gln Gln Pro Ala Val	
	710 715 720	
35	gcg ctg gaa acc tgc aac ccc cag ccc tgc cct gcc agg tgg gag	2205
	Ala Leu Glu Thr Cys Asn Pro Gln Pro Cys Pro Ala Arg Trp Glu	
40	725 730 735	
	gtg tca gag ccc agc tca tgc aca tca gct ggt gga gca ggc ctg	2250
	Val Ser Glu Pro Ser Ser Cys Thr Ser Ala Gly Gly Ala Gly Leu	
45	740 745 750	
	gcc ttg gag aac gag acc tgt gtg cca ggg gca gat ggc ctg gag	2295
50	Ala Leu Glu Asn Glu Thr Cys Val Pro Gly Ala Asp Gly Leu Glu	
	755 760 765	
	gct cca gtg act gag ggg cct ggc tcc gla gat gag aag ctg cct	2340
55	Ala Pro Val Thr Glu Gly Pro Gly Ser Val Asp Glu Lys Leu Pro	

	770	775	780	
5	gcc cct gag ccc tgt gtc ggg atg tca tgt cct cca ggc tgg ggc			2385
	Ala Pro Glu Pro Cys Val Gly Met Ser Cys Pro Pro Gly Trp Gly			
	785	790	795	
10	cat ctg gat gcc acc tct gca ggg gag aag gct ccc tcc cca tgg			2430
	His Leu Asp Ala Thr Ser Ala Gly Glu Lys Ala Pro Ser Pro Trp			
15	800	805	810	
	ggc agc atc agg acg ggg gct caa gct gca cac gtg tgg acc cct			2475
	Gly Ser Ile Arg Thr Gly Ala Gln Ala Ala His Val Trp Thr Pro			
20	815	820	825	
	gcg gca ggg tgc tgc tcc gtc tcc tgc ggg cga ggt ctg atg gag			2520
25	Ala Ala Gly Ser Cys Ser Val Ser Cys Gly Arg Gly Leu Met Glu			
	830	835	840	
	ctg cgt ttc ctg tgc atg gac tct gcc ctc agg gtg cct gtc cag			2565
30	Leu Arg Phe Leu Cys Met Asp Ser Ala Leu Arg Val Pro Val Gln			
	845	850	855	
35	gaa gag ctg tgt ggc ctg gca agc aag cct ggg agc cgg cgg gag			2610
	Glu Glu Leu Cys Gly Leu Ala Ser Lys Pro Gly Ser Arg Arg Glu			
	860	865	870	
40	gtc tgc cag gct gtc ccg tgc cct gct cgg tgg cag tac aag ctg			2655
	Val Cys Gln Ala Val Pro Cys Pro Ala Arg Trp Gln Tyr Lys Leu			
	875	880	885	
45	gcg gcc tgc agc gtg agc tgt ggg aga ggg gtc gtg cgg agg atc			2700
	Ala Ala Cys Ser Val Ser Cys Gly Arg Gly Val Val Arg Arg Ile			
50	890	895	900	
	ctg tat tgt gcc cgg gcc cat ggg gag gac gat ggt gag gag atc			2745
	Leu Tyr Cys Ala Arg Ala His Gly Glu Asp Asp Gly Glu Glu Ile			
55	905	910	915	

	ctg ttg gac acc cag tgc cag ggg ctg cct cgc ccg gaa ccc cag	2790
5	Leu Leu Asp Thr Gln Cys Gln Gly Leu Pro Arg Pro Glu Pro Gln	
	920 925 930	
	gag gcc tgc agc ctg gag ccc tgc cca cct agg tgg aaa gtc atg	2835
10	Glu Ala Cys Ser Leu Glu Pro Cys Pro Pro Arg Trp Lys Val Met	
	935 940 945	
15	tcc ctt ggc cca tgt tgc gcc agc tgt ggc ctt ggc act gct aga	2880
	Ser Leu Gly Pro Cys Ser Ala Ser Cys Gly Leu Gly Thr Ala Arg	
	950 955 960	
20	cgc tgc gtg gcc tgt gtg cag ctc gac caa ggc cag gac gtg gag	2925
	Arg Ser Val Ala Cys Val Gln Leu Asp Gln Gly Gln Asp Val Glu	
	965 970 975	
25	gtg gac gag gcg gcc tgt gcg gcg ctg gtg cgg ccc gag gcc agt	2970
	Val Asp Glu Ala Ala Cys Ala Ala Leu Val Arg Pro Glu Ala Ser	
30	980 985 990	
	gtc ccc tgt ctc att gcc gac tgc acc tac cgc tgg cat gtt ggc	3015
	Val Pro Cys Leu Ile Ala Asp Cys Thr Tyr Arg Trp His Val Gly	
35	995 1000 1005	
	acc tgg atg gag tgc tct gtt tcc tgt ggg gat ggc atc cag cgc	3060
40	Thr Trp Met Glu Cys Ser Val Ser Cys Gly Asp Gly Ile Gln Arg	
	1010 1015 1020	
	cgg cgt gac acc tgc ctc gga ccc cag gcc cag gcg cct gtg cca	3105
45	Arg Arg Asp Thr Cys Leu Gly Pro Gln Ala Gln Ala Pro Val Pro	
	1025 1030 1035	
50	gct gat ttc tgc cag cac ttg ccc aag ccg gtg act gtg cgt ggc	3150
	Ala Asp Phe Cys Gln His Leu Pro Lys Pro Val Thr Val Arg Gly	
	1040 1045 1050	
55	tgc tgg gct ggg ccc tgt gtg gga cag ggt gcc tgt ggc agg cag	3195

	Cys Trp Ala Gly Pro Cys Val Gly Gln Gly Ala Cys Gly Arg Gln	
5	1055 1060 1065	
	cac ctt gag cca aca gga acc att gac atg cga ggc cca ggg cag	3240
	His Leu Glu Pro Thr Gly Thr Ile Asp Met Arg Gly Pro Gly Gln	
10	1070 1075 1080	
	gca gac tgt gca gtg gcc att ggg cgg ccc ctc ggg gag gtg gtg	3285
15	Ala Asp Cys Ala Val Ala Ile Gly Arg Pro Leu Gly Glu Val Val	
	1085 1090 1095	
	acc ctc cgc gtc ctt gag agt tct ctc aac tgc agt gcg ggg gac	3330
20	Thr Leu Arg Val Leu Glu Ser Ser Leu Asn Cys Ser Ala Gly Asp	
	1100 1105 1110	
25	atg ttg ctg ctt tgg ggc cgg ctc acc tgg agg aag atg tgc agg	3375
	Met Leu Leu Leu Trp Gly Arg Leu Thr Trp Arg Lys Met Cys Arg	
	1115 1120 1125	
30	aag ctg ttg gac atg act ttc agc tcc aag acc aac acg ctg gtg	3420
	Lys Leu Leu Asp Met Thr Phe Ser Ser Lys Thr Asn Thr Leu Val	
	1130 1135 1140	
35	gtg agg cag cgc tgc ggg cgg cca gga ggt ggg gtg ctg ctg cgg	3465
	Val Arg Gln Arg Cys Gly Arg Pro Gly Gly Gly Val Leu Leu Arg	
40	1145 1150 1155	
	tat ggg agc cag ctt gct cct gaa acc ttc tac aga gaa tgt gac	3510
	Tyr Gly Ser Gln Leu Ala Pro Glu Thr Phe Tyr Arg Glu Cys Asp	
45	1160 1165 1170	
	atg cag ctc ttt ggg ccc tgg ggt gaa atc gtg agc ccc tcg ctg	3555
50	Met Gln Leu Phe Gly Pro Trp Gly Glu Ile Val Ser Pro Ser Leu	
	1175 1180 1185	
	agt cca gcc acg agt aat gca ggg ggc tgc cgg ctc ttc att aat	3600
55	Ser Pro Ala Thr Ser Asn Ala Gly Gly Cys Arg Leu Phe Ile Asn	

	1190	1195	1200	
5	gtg gct ccg cac gca cgg att gcc atc cat gcc ctg gcc acc aac			3645
	Val Ala Pro His Ala Arg Ile Ala Ile His Ala Leu Ala Thr Asn			
	1205	1210	1215	
10	atg ggc gct ggg acc gag gga gcc aat gcc agc tac atc ttg atc			3690
	Met Gly Ala Gly Thr Glu Gly Ala Asn Ala Ser Tyr Ile Leu Ile			
	1220	1225	1230	
15	cgg gac acc cac agc ttg agg acc aca gcg ttc cat ggg cag cag			3735
	Arg Asp Thr His Ser Leu Arg Thr Thr Ala Phe His Gly Gln Gln			
20	1235	1240	1245	
	gtg ctc tac tgg gag tca gag agc agc cag gct gag atg gag ttc			3780
25	Val Leu Tyr Trp Glu Ser Glu Ser Ser Gln Ala Glu Met Glu Phe			
	1250	1255	1260	
30	agc gag ggc ttc ctg aag gct cag gcc agc ctg cgg ggc cag tac			3825
	Ser Glu Gly Phe Leu Lys Ala Gln Ala Ser Leu Arg Gly Gln Tyr			
	1265	1270	1275	
35	tgg acc ctc caa tca tgg gta ccg gag atg cag gac cct cag tcc			3870
	Trp Thr Leu Gln Ser Trp Val Pro Glu Met Gln Asp Pro Gln Ser			
	1280	1285	1290	
40	tgg aag gga aag gaa gga acc			3891
	Trp Lys Gly Lys Glu Gly Thr			
	1295			
45				
	<210>18			
50	<211>1378			
	<212>PRT			
	<213> Homo sapiens			
55	<400>18			

	gct gca ggc ggc atc cta cac ctg gag ctg ctg gtg gcc gtg ggc	45
5	Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly	
	1 5 10 15	
10	ccc gat gtc ttc cag gct cac cag gag gac aca gag cgc tat gtg	90
	Pro Asp Val Phe Gln Ala His Gln Glu Asp Thr Glu Arg Tyr Val	
	20 25 30	
15	ctc acc aac ctc aac atc ggg gca gaa ctg ctt cgg gac ccg tcc	135
	Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro Ser	
	35 40 45	
20	ctg ggg gct cag ttt cgg gtg cac ctg gtg aag atg gtc att ctg	180
	Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile Leu	
	50 55 60	
25	aca gag cct gag ggt gct cca aat atc aca gcc aac ctc acc tcg	225
	Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr Ser	
30	65 70 75	
	tcc ctg ctg agc gtc tgt ggg tgg agc cag acc atc aac cct gag	270
	Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro Glu	
35	80 85 90	
	gac gac acg gat cct ggc cat gct gac ctg gtc ctc tat atc act	315
40	Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile Thr	
	95 100 105	
45	agg ttt gac ctg gag ttg cct gat ggt aac cgg cag gtg cgg ggc	360
	Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg Gly	
	110 115 120	
50	gtc acc cag ctg ggc ggt gcc tgc tcc cca acc tgg agc tgc ctc	405
	Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys Leu	
	125 130 135	
55	att acc gag gac act ggc ttc gac ctg gga gtc acc att gcc cat	450

	Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala His	
5	140 145 150	
	gag att ggc cac agc ttc ggc ctg gag cac gac ggc gcg ccc ggc	495
	Glu Ile Gly His Ser Phe Gly Leu Glu His Asp Gly Ala Pro Gly	
10	155 160 165	
	agc ggc tgc ggc ccc agc gga cac gtg atg gct tgc gac ggc gcc	540
15	Ser Gly Cys Gly Pro Ser Gly His Val Met Ala Ser Asp Gly Ala	
	170 175 180	
	gcg ccc cgc gcc ggc ctc gcc tgg tcc ccc tgc agc cgc cgg cag	585
20	Ala Pro Arg Ala Gly Leu Ala Trp Ser Pro Cys Ser Arg Arg Gln	
	185 190 195	
25	ctg ctg agc ctg ctc agg acg ggc gcg ctg cgt gtg gga ccc gcc	630
	Leu Leu Ser Leu Leu Arg Thr Gly Ala Leu Arg Val Gly Pro Ala	
	200 205 210	
30	gcg gcc tca acc cgg gtc cgc ggg gca ccc gcc gga tgc gca gcc	675
	Ala Ala Ser Thr Arg Val Arg Gly Ala Pro Ala Gly Cys Ala Ala	
	215 220 225	
35	tgg cct cta cta cag cgc caa cga gca gtg ccg cgt ggc ctt cgg	720
	Trp Pro Leu Leu Gln Arg Gln Arg Ala Val Pro Arg Gly Leu Arg	
40	230 235 240	
	ccc caa ggc tgt cgc ctg cac ctt cgc cag gga gca cct ggt gag	765
	Pro Gln Gly Cys Arg Leu His Leu Arg Gln Gly Ala Pro Gly Glu	
45	245 250 255	
	tct gcc ggc ggt ggc ctg gga ttg gct gtg agg tcc ctc cgc atc	810
50	Ser Ala Gly Gly Gly Leu Gly Leu Ala Val Arg Ser Leu Arg Ile	
	260 265 270	
	acc cag ctc acg tcc ccc caa acg tgc atg gat atg tgc cag gcc	855
55	Thr Gln Leu Thr Ser Pro Gln Thr Cys Met Asp Met Cys Gln Ala	

	275	280	285	
5	ctc tcc tgc cac aca gac ccg ctg gac caa agc agc tgc agc cgc			900
	Leu Ser Cys His Thr Asp Pro Leu Asp Gln Ser Ser Cys Ser Arg			
	290	295	300	
10	ctc ctc gtt cct ctc ctg gat ggg aca gaa tgt ggc gtg gag aag			945
	Leu Leu Val Pro Leu Leu Asp Gly Thr Glu Cys Gly Val Glu Lys			
	305	310	315	
15	tgg tgc tcc aag ggt cgc tgc cgc tcc ctg gtg gag ctg acc ccc			990
	Trp Cys Ser Lys Gly Arg Cys Arg Ser Leu Val Glu Leu Thr Pro			
20	320	325	330	
	ata gca gca gtg cat ggg cgc tgg tct agc tgg ggt ccc cga agt			1035
	Ile Ala Ala Val His Gly Arg Trp Ser Ser Trp Gly Pro Arg Ser			
25	335	340	345	
	cct tgc tcc cgc tcc tgc gga gga ggt gtg gtc acc agg agg cgg			1080
30	Pro Cys Ser Arg Ser Cys Gly Gly Gly Val Val Thr Arg Arg Arg			
	350	355	360	
35	cag tgc aac aac ccc aga cct gcc ttt ggg ggg cgt gca tgt gtt			1125
	Gln Cys Asn Asn Pro Arg Pro Ala Phe Gly Gly Arg Ala Cys Val			
	365	370	375	
40	ggt gct gac ctc cag gcc gag atg tgc aac act cag gcc tgc gag			1170
	Gly Ala Asp Leu Gln Ala Glu Met Cys Asn Thr Gln Ala Cys Glu			
	380	385	390	
45	aag acc cag ctg gag ttc atg tcg caa cag tgc gcc agg acc gac			1215
	Lys Thr Gln Leu Glu Phe Met Ser Gln Gln Cys Ala Arg Thr Asp			
	395	400	405	
50	ggc cag ccg ctg cgc tcc tcc cct ggc ggc gcc tcc ttc tac cac			1260
	Gly Gln Pro Leu Arg Ser Ser Pro Gly Gly Ala Ser Phe Tyr His			
55	410	415	420	

	tgg ggt gct gct gta cca cac agc caa ggg gat gct ctg tgc aga	1305
5	Trp Gly Ala Ala Val Pro His Ser Gln Gly Asp Ala Leu Cys Arg	
	425 430 435	
	cac atg tgc cgg gcc att ggc gag agc ttc atc atg aag cgt gga	1350
10	His Met Cys Arg Ala Ile Gly Glu Ser Phe Ile Met Lys Arg Gly	
	440 445 450	
15	gac agc ttc ctc gat ggg acc cgg tgt atg cca agt ggc ccc cgg	1395
	Asp Ser Phe Leu Asp Gly Thr Arg Cys Met Pro Ser Gly Pro Arg	
	455 460 465	
20	gag gac ggg acc ctg agc ctg tgt gtg tgc ggc agc tgc agg aca	1440
	Glu Asp Gly Thr Leu Ser Leu Cys Val Ser Gly Ser Cys Arg Thr	
	470 475 480	
25	ttt ggc tgt gat ggt agg atg gac tcc cag cag gta tgg gac agg	1485
	Phe Gly Cys Asp Gly Arg Met Asp Ser Gln Gln Val Trp Asp Arg	
30	485 490 495	
	tgc cag gtg tgt ggt ggg gac aac agc acg tgc agc cca cgg aag	1530
	Cys Gln Val Cys Gly Gly Asp Asn Ser Thr Cys Ser Pro Arg Lys	
35	500 505 510	
	ggc tct ttc aca gct ggc aga gcg aga gaa tat gtc acg ttt ctg	1575
40	Gly Ser Phe Thr Ala Gly Arg Ala Arg Glu Tyr Val Thr Phe Leu	
	515 520 525	
	aca gtt acc ccc aac ctg acc agt gtc tac att gcc aac cac agg	1620
45	Thr Val Thr Pro Asn Leu Thr Ser Val Tyr Ile Ala Asn His Arg	
	530 535 540	
50	cct ctc ttc aca cac ttg gcg gtg agg atc gga ggg cgc tat gtc	1665
	Pro Leu Phe Thr His Leu Ala Val Arg Ile Gly Gly Arg Tyr Val	
	545 550 555	
55	gtg gct ggg aag atg agc atc tcc cct aac acc acc tac ccc tcc	1710

	Val Ala Gly Lys Met Ser Ile Ser Pro Asn Thr Thr Tyr Pro Ser	
5	560 565 570	
	ctc ctg gag gat ggt cgt gtc gag tac aga gtg gcc ctc acc gag	1755
10	Leu Leu Glu Asp Gly Arg Val Glu Tyr Arg Val Ala Leu Thr Glu	
	575 580 585	
	gac cgg ctg ccc cgc ctg gag gag atc cgc atc tgg gga ccc ctc	1800
15	Asp Arg Leu Pro Arg Leu Glu Glu Ile Arg Ile Trp Gly Pro Leu	
	590 595 600	
20	cag gaa gat gct gac atc cag gtt tac agg cgg tat ggc gag gag	1845
	Gln Glu Asp Ala Asp Ile Gln Val Tyr Arg Arg Tyr Gly Glu Glu	
	605 610 615	
25	tat ggc aac ctc acc cgc cca gac atc acc ttc acc tac ttc cag	1890
	Tyr Gly Asn Leu Thr Arg Pro Asp Ile Thr Phe Thr Tyr Phe Gln	
	620 625 630	
30	cct aag cca cgg cag gcc tgg gtg tgg gcc gct gtg cgt ggg ccc	1935
	Pro Lys Pro Arg Gln Ala Trp Val Trp Ala Ala Val Arg Gly Pro	
	635 640 645	
35	tgc tgc gtg agc tgt ggg gca ggg ctg cgc tgg gta aac tac agc	1980
	Cys Ser Val Ser Cys Gly Ala Gly Leu Arg Trp Val Asn Tyr Ser	
40	650 655 660	
	tgc ctg gac cag gcc agg aag gag ttg gtg gag act gtc cag tgc	2025
	Cys Leu Asp Gln Ala Arg Lys Glu Leu Val Glu Thr Val Gln Cys	
45	665 670 675	
	caa ggg agc cag cag cca cca gcg tgg cca gag gcc tgc gtg ctc	2070
50	Gln Gly Ser Gln Gln Pro Pro Ala Trp Pro Glu Ala Cys Val Leu	
	680 685 690	
	gaa ccc tgc cct ccc tac tgg gcg gtg gga gac ttc ggc cca tgc	2115
55	Glu Pro Cys Pro Pro Tyr Trp Ala Val Gly Asp Phe Gly Pro Cys	

	695	700	705	
5	agc gcc tcc tgt ggg ggc ggc ctg cgg gag cgg cca gtg cgc tgc			2160
	Ser Ala Ser Cys Gly Gly Gly Leu Arg Glu Arg Pro Val Arg Cys			
	710	715	720	
10	gtg gag gcc cag ggc agc ctc ctg aag aca ttg ccc cca gcc cgg			2205
	Val Glu Ala Gln Gly Ser Leu Leu Lys Thr Leu Pro Pro Ala Arg			
	725	730	735	
15	tgc aga gca ggg gcc cag cag cca gct gtg gcg ctg gaa acc tgc			2250
	Cys Arg Ala Gly Ala Gln Gln Pro Ala Val Ala Leu Glu Thr Cys			
	740	745	750	
20	aac ccc cag ccc tgc cct gcc agg tgg gag gtg tca gag ccc agc			2295
	Asn Pro Gln Pro Cys Pro Ala Arg Trp Glu Val Ser Glu Pro Ser			
25	755	760	765	
	tca tgc aca tca gct ggt gga gca ggc ctg gcc ttg gag aac gag			2340
30	Ser Cys Thr Ser Ala Gly Gly Ala Gly Leu Ala Leu Glu Asn Glu			
	770	775	780	
35	acc tgt gtg cca ggg gca gat ggc ctg gag gct cca gtg act gag			2385
	Thr Cys Val Pro Gly Ala Asp Gly Leu Glu Ala Pro Val Thr Glu			
	785	790	795	
40	ggg cct ggc tcc gla gat gag aag ctg cct gcc cct gag ccc tgt			2430
	Gly Pro Gly Ser Val Asp Glu Lys Leu Pro Ala Pro Glu Pro Cys			
	800	805	810	
45	gtc ggg atg tca tgt cct cca ggc tgg ggc cat ctg gat gcc acc			2475
	Val Gly Met Ser Cys Pro Pro Gly Trp Gly His Leu Asp Ala Thr			
	815	820	825	
50	tct gca ggg gag aag gct ccc tcc cca tgg ggc agc atc agg acg			2520
	Ser Ala Gly Glu Lys Ala Pro Ser Pro Trp Gly Ser Ile Arg Thr			
55	830	835	840	

	ggg gct caa gct gca cac gtg tgg acc cct gcg gca ggg tgc tgc	2565
5	Gly Ala Gln Ala Ala His Val Trp Thr Pro Ala Ala Gly Ser Cys	
	845 850 855	
	tcc gtc tcc tgc ggg cga ggt ctg atg gag ctg cgt ttc ctg tgc	2610
10	Ser Val Ser Cys Gly Arg Gly Leu Met Glu Leu Arg Phe Leu Cys	
	860 865 870	
	atg gac tct gcc ctc agg gtg cct gtc cag gaa gag ctg tgt ggc	2655
15	Met Asp Ser Ala Leu Arg Val Pro Val Gln Glu Glu Leu Cys Gly	
	875 880 885	
	ctg gca agc aag cct ggg agc cgg cgg gag gtc tgc cag gct gtc	2700
20	Leu Ala Ser Lys Pro Gly Ser Arg Arg Glu Val Cys Gln Ala Val	
	890 895 900	
	ccg tgc cct gct cgg tgg cag tac aag ctg gcg gcc tgc agc gtg	2745
25	Pro Cys Pro Ala Arg Trp Gln Tyr Lys Leu Ala Ala Cys Ser Val	
	905 910 915	
	agc tgt ggg aga ggg gtc gtg cgg agg atc ctg tat tgt gcc cgg	2790
30	Ser Cys Gly Arg Gly Val Val Arg Arg Ile Leu Tyr Cys Ala Arg	
	920 925 930	
	gcc cat ggg gag gac gat ggt gag gag atc ctg ttg gac acc cag	2835
35	Ala His Gly Glu Asp Asp Gly Glu Glu Ile Leu Leu Asp Thr Gln	
	935 940 945	
	tgc cag ggg ctg cct cgc ccg gaa ccc cag gag gcc tgc agc ctg	2880
40	Cys Gln Gly Leu Pro Arg Pro Glu Pro Gln Glu Ala Cys Ser Leu	
	950 955 960	
	gag ccc tgc cca cct agg tgg aaa gtc atg tcc ctt ggc cca tgt	2925
45	Glu Pro Cys Pro Pro Arg Trp Lys Val Met Ser Leu Gly Pro Cys	
	965 970 975	
	tgc gcc agc tgt ggc ctt ggc act gct aga cgc tgc gtg gcc tgt	2970
50		
55		

	Ser Ala Ser Cys Gly Leu Gly Thr Ala Arg Arg Ser Val Ala Cys	
5	980 985 990	
	gtg cag ctc gac caa ggc cag gac gtg gag gtg gac gag gcg gcc	3015
	Val Gln Leu Asp Gln Gly Gln Asp Val Glu Val Asp Glu Ala Ala	
10	995 1000 1005	
	tgt gcg gcg ctg gtg cgg ccc gag gcc agt gtc ccc tgt ctc att	3060
15	Cys Ala Ala Leu Val Arg Pro Glu Ala Ser Val Pro Cys Leu Ile	
	1010 1015 1020	
	gcc gac tgc acc tac cgc tgg cat gtt ggc acc tgg atg gag tgc	3105
20	Ala Asp Cys Thr Tyr Arg Trp His Val Gly Thr Trp Met Glu Cys	
	1025 1030 1035	
25	tct gtt tcc tgt ggg gat ggc atc cag cgc cgg cgt gac acc tgc	3150
	Ser Val Ser Cys Gly Asp Gly Ile Gln Arg Arg Arg Asp Thr Cys	
	1040 1045 1050	
30	ctc gga ccc cag gcc cag gcg cct gtg cca gct gat ttc tgc cag	3195
	Leu Gly Pro Gln Ala Gln Ala Pro Val Pro Ala Asp Phe Cys Gln	
	1055 1060 1065	
35	cac ttg ccc aag ccg gtg act gtg cgt ggc tgc tgg gct ggg ccc	3240
	His Leu Pro Lys Pro Val Thr Val Arg Gly Cys Trp Ala Gly Pro	
40	1070 1075 1080	
	tgt gtg gga cag ggt acg ccc agc ctg gtg ccc cac gaa gaa gcc	3285
	Cys Val Gly Gln Gly Thr Pro Ser Leu Val Pro His Glu Glu Ala	
45	1085 1090 1095	
	gct gct cca gga cgg acc aca gcc acc cct gct ggt gcc tcc ctg	3330
50	Ala Ala Pro Gly Arg Thr Thr Ala Thr Pro Ala Gly Ala Ser Leu	
	1100 1105 1110	
	gag tgg tcc cag gcc cgg ggc ctg ctc ttc tcc ccg gct ccc cag	3375
55	Glu Trp Ser Gln Ala Arg Gly Leu Leu Phe Ser Pro Ala Pro Gln	

	1115	1120	1125	
5	cct cgg cgg ctc ctg ccc ggg ccc cag gaa aac tca gtg cag tcc			3420
	Pro Arg Arg Leu Leu Pro Gly Pro Gln Glu Asn Ser Val Gln Ser			
	1130	1135	1140	
10	agt gcc tgt ggc agg cag cac ctt gag cca aca gga acc att gac			3465
	Ser Ala Cys Gly Arg Gln His Leu Glu Pro Thr Gly Thr Ile Asp			
	1145	1150	1155	
15	atg cga ggc cca ggg cag gca gac tgt gca gtg gcc att ggg cgg			3510
	Met Arg Gly Pro Gly Gln Ala Asp Cys Ala Val Ala Ile Gly Arg			
20	1160	1165	1170	
	ccc ctc ggg gag gtg gtg acc ctc cgc gtc ctt gag agt tct ctc			3555
25	Pro Leu Gly Glu Val Val Thr Leu Arg Val Leu Glu Ser Ser Leu			
	1175	1180	1185	
	aac tgc agt gcg ggg gac atg ttg ctg ctt tgg ggc cgg ctc acc			3600
30	Asn Cys Ser Ala Gly Asp Met Leu Leu Leu Trp Gly Arg Leu Thr			
	1190	1195	1200	
35	tgg agg aag atg tgc agg aag ctg ttg gac atg act ttc agc tcc			3645
	Trp Arg Lys Met Cys Arg Lys Leu Leu Asp Met Thr Phe Ser Ser			
	1205	1210	1215	
40	aag acc aac acg ctg gtg gtg agg cag cgc tgc ggg cgg cca gga			3690
	Lys Thr Asn Thr Leu Val Val Arg Gln Arg Cys Gly Arg Pro Gly			
	1220	1225	1230	
45	ggg ggg gtg ctg ctg cgg tat ggg agc cag ctt gct cct gaa acc			3735
	Gly Gly Val Leu Leu Arg Tyr Gly Ser Gln Leu Ala Pro Glu Thr			
	1235	1240	1245	
50	ttc tac aga gaa tgt gac atg cag ctc ttt ggg ccc tgg ggt gaa			3780
	Phe Tyr Arg Glu Cys Asp Met Gln Leu Phe Gly Pro Trp Gly Glu			
55	1250	1255	1260	

	atc gtg agc ccc tgc ctg agt cca gcc acg agt aat gca ggg ggc	3825
5	Ile Val Ser Pro Ser Leu Ser Pro Ala Thr Ser Asn Ala Gly Gly	
	1265 1270 1275	
10	tgc cgg ctc ttc att aat gtg gct ccg cac gca cgg att gcc atc	3870
	Cys Arg Leu Phe Ile Asn Val Ala Pro His Ala Arg Ile Ala Ile	
	1280 1285 1290	
15	cat gcc ctg gcc acc aac atg ggc gct ggg acc gag gga gcc aat	3915
	His Ala Leu Ala Thr Asn Met Gly Ala Gly Thr Glu Gly Ala Asn	
	1295 1300 1305	
20	gcc agc tac atc ttg atc cgg gac acc cac agc ttg agg acc aca	3960
	Ala Ser Tyr Ile Leu Ile Arg Asp Thr His Ser Leu Arg Thr Thr	
	1310 1315 1320	
25	gcg ttc cat ggg cag cag gtg ctc tac tgg gag tca gag agc agc	4005
	Ala Phe His Gly Gln Gln Val Leu Tyr Trp Gly Ser Glu Ser Ser	
30	1325 1330 1335	
	cag gct gag atg gag ttc agc gag ggc ttc ctg aag gct cag gcc	4050
	Gln Ala Glu Met Glu Phe Ser Glu Gly Phe Leu Lys Ala Gln Ala	
35	1340 1345 1350	
	agc ctg cgg ggc cag tac tgg acc ctc caa tca tgg gta ccg gag	4095
40	Ser Leu Arg Gly Gln Tyr Trp Thr Leu Gln Ser Trp Val Pro Glu	
	1355 1360 1365	
45	atg cag gac cct cag tcc tgg aag gga aag gaa gga acc	4134
	Met Gln Asp Pro Gln Ser Trp Lys Gly Lys Glu Gly Thr	
	1370 1375	
50	<210>19	
	<211>1322	
55	<212>PRT	

<213> Homo sapiens

<400>19

5	gct gca ggc ggc atc cta cac ctg gag ctg ctg gtg gcc gtg ggc	45
	Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly	
10	1 5 10 15	
	ccc gat gtc ttc cag gct cac cag gag gac aca gag cgc tat gtg	90
15	Pro Asp Val Phe Gln Ala His Gln Glu Asp Thr Glu Arg Tyr Val	
	20 25 30	
20	ctc acc aac ctc aac atc ggg gca gaa ctg ctt cgg gac ccg tcc	135
	Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro Ser	
	35 40 45	
25	ctg ggg gct cag ttt cgg gtg cac ctg gtg aag atg gtc att ctg	180
	Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile Leu	
	50 55 60	
30	aca gag cct gag ggt gct cca aat atc aca gcc aac ctc acc tgc	225
	Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr Ser	
	65 70 75	
35	tcc ctg ctg agc gtc tgt ggg tgg agc cag acc atc aac cct gag	270
	Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro Glu	
40	80 85 90	
	gac gac acg gat cct ggc cat gct gac ctg gtc ctc tat atc act	315
	Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile Thr	
45	95 100 105	
	agg ttt gac ctg gag ttg cct gat ggt aac cgg cag gtg cgg ggc	360
50	Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg Gly	
	110 115 120	
55	gtc acc cag ctg ggc ggt gcc tgc tcc cca acc tgg agc tgc ctc	405
	Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys Leu	

	125	130	135	
5	att acc gag gac act ggc ttc gac ctg gga gtc acc att gcc cat			450
	Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala His			
	140	145	150	
10	gag att ggg cac agc ttc ggc ctg gag cac gac ggc gcg ccc ggc			495
	Glu Ile Gly His Ser Phe Gly Leu Glu His Asp Gly Ala Pro Gly			
	155	160	165	
15	agc ggc tgc ggc ccc agc gga cac gtg atg gct tgc gac ggc gcc			540
	Ser Gly Cys Gly Pro Ser Gly His Val Met Ala Ser Asp Gly Ala			
20	170	175	180	
	gcg ccc cgc gcc ggc ctc gcc tgg tcc ccc tgc agc cgc cgg cag			585
25	Ala Pro Arg Ala Gly Leu Ala Trp Ser Pro Cys Ser Arg Arg Gln			
	185	190	195	
	ctg ctg agc ctg ctc agg acg ggc gcg ctg cgt gtg gga ccc gcc			630
30	Leu Leu Ser Leu Leu Arg Thr Gly Ala Leu Arg Val Gly Pro Ala			
	200	205	210	
	gcg gcc tca acc cgg gtc cgc ggg gca ccc gcc gga tgc gca gcc			675
35	Ala Ala Ser Thr Arg Val Arg Gly Ala Pro Ala Gly Cys Ala Ala			
	215	220	225	
40	tgg cct cta cta cag cgc caa cga gca gtg ccg cgt ggc ctt cgg			720
	Trp Pro Leu Leu Gln Arg Gln Arg Ala Val Pro Arg Gly Leu Arg			
	230	235	240	
45	ccc caa ggc tgt cgc ctg cac ctt cgc cag gga gca cct ggt gag			765
	Pro Gln Gly Cys Arg Leu His Leu Arg Gln Gly Ala Pro Gly Glu			
	245	250	255	
50	tct gcc ggc ggt ggc ctg gga ttg gct gtg agg tcc ctc cgc atc			810
	Ser Ala Gly Gly Gly Leu Gly Leu Ala Val Arg Ser Leu Arg Ile			
55	260	265	270	

	acc cag ctc acg tcc ccc caa acg tgc atg gat atg tgc cag gcc	855
5	Thr Gln Leu Thr Ser Pro Gln Thr Cys Met Asp Met Cys Gln Ala	
	275 280 285	
	ctc tcc tgc cac aca gac ccg ctg gac caa agc agc tgc agc cgc	900
10	Leu Ser Cys His Thr Asp Pro Leu Asp Gln Ser Ser Cys Ser Arg	
	290 295 300	
	ctc ctc gtt cct ctc ctg gat ggg aca gaa tgt ggc gtg gag aag	945
15	Leu Leu Val Pro Leu Leu Asp Gly Thr Glu Cys Gly Val Glu Lys	
	305 310 315	
20	tgg tgc tcc aag ggt cgc tgc cgc tcc ctg gtg gag ctg acc ccc	990
	Trp Cys Ser Lys Gly Arg Cys Arg Ser Leu Val Glu Leu Thr Pro	
	320 325 330	
25	ata gca gca gtg cat ggg cgc tgg tct agc tgg ggt ccc cga agt	1035
	Ile Ala Ala Val His Gly Arg Trp Ser Ser Trp Gly Pro Arg Ser	
30	335 340 345	
	cct tgc tcc cgc tcc tgc gga gga ggt gtg gtc acc agg agg cgc	1080
	Pro Cys Ser Arg Ser Cys Gly Gly Gly Val Val Thr Arg Arg Arg	
35	350 355 360	
	cag tgc aac aac ccc aga cct gcc ttt ggg ggg cgt gca tgt gtt	1125
40	Gln Cys Asn Asn Pro Arg Pro Ala Phe Gly Gly Arg Ala Cys Val	
	365 370 375	
	ggt gct gac ctc cag gcc gag atg tgc aac act cag gcc tgc gag	1170
45	Gly Ala Asp Leu Gln Ala Glu Met Cys Asn Thr Gln Ala Cys Glu	
	380 385 390	
	aag acc cag ctg gag ttc atg tgc caa cag tgc gcc agg acc gac	1215
50	Lys Thr Gln Leu Glu Phe Met Ser Gln Gln Cys Ala Arg Thr Asp	
	395 400 405	
55	ggc cag ccg ctg cgc tcc tcc cct ggc ggc gcc tcc ttc tac cac	1260

	Gly Gln Pro Leu Arg Ser Ser Pro Gly Gly Ala Ser Phe Tyr His	
5	410 415 420	
	tgg ggt gct gct gta cca cac agc caa ggg gat gct ctg tgc aga	1305
10	Trp Gly Ala Ala Val Pro His Ser Gln Gly Asp Ala Leu Cys Arg	
	425 430 435	
	cac atg tgc cgg gcc att ggc gag agc ttc atc atg aag cgt gga	1350
15	His Met Cys Arg Ala Ile Gly Glu Ser Phe Ile Met Lys Arg Gly	
	440 445 450	
20	gac agc ttc ctc gat ggg acc cgg tgt atg cca agt ggc ccc cgg	1395
	Asp Ser Phe Leu Asp Gly Thr Arg Cys Met Pro Ser Gly Pro Arg	
	455 460 465	
25	gag gac ggg acc ctg agc ctg tgt gtg tgc ggc agc tgc agg aca	1440
	Glu Asp Gly Thr Leu Ser Leu Cys Val Ser Gly Ser Cys Arg Thr	
	470 475 480	
30	ttt ggc tgt gat ggt agg atg gac tcc cag cag gta tgg gac agg	1485
	Phe Gly Cys Asp Gly Arg Met Asp Ser Gln Gln Val Trp Asp Arg	
	485 490 495	
35	tgc cag gtg tgt ggt ggg gac aac agc acg tgc agc cca cgg aag	1530
	Cys Gln Val Cys Gly Gly Asp Asn Ser Thr Cys Ser Pro Arg Lys	
40	500 505 510	
	ggc tct ttc aca gct ggc aga gcg aga gaa tat gtc acg ttt ctg	1575
45	Gly Ser Phe Thr Ala Gly Arg Ala Arg Glu Tyr Val Thr Phe Leu	
	515 520 525	
	aca gtt acc ccc aac ctg acc agt gtc tac att gcc aac cac agg	1620
50	Thr Val Thr Pro Asn Leu Thr Ser Val Tyr Ile Ala Asn His Arg	
	530 535 540	
55	cct ctc ttc aca cac ttg gcg gtg agg atc gga ggg cgc tat gtc	1665
	Pro Leu Phe Thr His Leu Ala Val Arg Ile Gly Gly Arg Tyr Val	

	545	550	555	
5	gtg gct ggg aag atg agc atc tcc cct aac acc acc tac ccc tcc			1710
	Val Ala Gly Lys Met Ser Ile Ser Pro Asn Thr Thr Tyr Pro Ser			
	560	565	570	
10	ctc ctg gag gat ggt cgt gtc gag tac aga gtg gcc ctc acc gag			1755
	Leu Leu Glu Asp Gly Arg Val Glu Tyr Arg Val Ala Leu Thr Glu			
15	575	580	585	
	gac cgg ctg ccc cgc ctg gag gag atc cgc atc tgg gga ccc ctc			1800
	Asp Arg Leu Pro Arg Leu Glu Glu Ile Arg Ile Trp Gly Pro Leu			
20	590	595	600	
	cag gaa gat gct gac atc cag gtt tac agg cgg tat ggc gag gag			1845
25	Gln Glu Asp Ala Asp Ile Gln Val Tyr Arg Arg Tyr Gly Glu Glu			
	605	610	615	
	tat ggc aac ctc acc cgc cca gac atc acc ttc acc tac ttc cag			1890
30	Tyr Gly Asn Leu Thr Arg Pro Asp Ile Thr Phe Thr Tyr Phe Gln			
	620	625	630	
	cct aag cca cgg cag gcc tgg gtg tgg gcc gct gtg cgt ggg ccc			1935
35	Pro Lys Pro Arg Gln Ala Trp Val Trp Ala Ala Val Arg Gly Pro			
	635	640	645	
40	tgc tcg gtg agc tgt ggg gca ggg ctg cgc tgg gta aac tac agc			1980
	Cys Ser Val Ser Cys Gly Ala Gly Leu Arg Trp Val Asn Tyr Ser			
	650	655	660	
45	tgc ctg gac cag gcc agg aag gag ttg gtg gag act gtc cag tgc			2025
	Cys Leu Asp Gln Ala Arg Lys Glu Leu Val Glu Thr Val Gln Cys			
	665	670	675	
50	caa ggg agc cag cag cca cca gcg tgg cca gag gcc tgc gtg ctc			2070
	Gln Gly Ser Gln Gln Pro Pro Ala Trp Pro Glu Ala Cys Val Leu			
55	680	685	690	

	gaa ccc tgc cct ccc tac tgg gcg gtg gga gac ttc ggc cca tgc	2115
5	Glu Pro Cys Pro Pro Tyr Trp Ala Val Gly Asp Phe Gly Pro Cys	
	695 700 705	
	agc gcc tcc tgt ggg ggc ggc ctg cgg gag cgg cca gtg cgc tgc	2160
10	Ser Ala Ser Cys Gly Gly Gly Leu Arg Glu Arg Pro Val Arg Cys	
	710 715 720	
	gtg gag gcc cag ggc agc ctc ctg aag aca ttg ccc cca gcc cgg	2205
15	Val Glu Ala Gln Gly Ser Leu Leu Lys Thr Leu Pro Pro Ala Arg	
	725 730 735	
	tgc aga gca ggg gcc cag cag cca gct gtg gcg ctg gaa acc tgc	2250
20	Cys Arg Ala Gly Ala Gln Gln Pro Ala Val Ala Leu Glu Thr Cys	
	740 745 750	
	aac ccc cag ccc tgc cct gcc agg tgg gag gtg tca gag ccc agc	2295
25	Asn Pro Gln Pro Cys Pro Ala Arg Trp Glu Val Ser Glu Pro Ser	
	755 760 765	
	tca tgc aca tca gct ggt gga gca ggc ctg gcc ttg gag aac gag	2340
30	Ser Cys Thr Ser Ala Gly Gly Ala Gly Leu Ala Leu Glu Asn Glu	
	770 775 780	
	acc tgt gtg cca ggg gca gat ggc ctg gag gct cca gtg act gag	2385
35	Thr Cys Val Pro Gly Ala Asp Gly Leu Glu Ala Pro Val Thr Glu	
	785 790 795	
	ggg cct ggc tcc gta gat gag aag ctg cct gcc cct gag ccc tgt	2430
40	Gly Pro Gly Ser Val Asp Glu Lys Leu Pro Ala Pro Glu Pro Cys	
	800 805 810	
	gtc ggg atg tca tgt cct cca ggc tgg ggc cat ctg gat gcc acc	2475
45	Val Gly Met Ser Cys Pro Pro Gly Trp Gly His Leu Asp Ala Thr	
	815 820 825	
	tct gca ggg gag aag gct ccc tcc cca tgg ggc agc atc agg acg	2520
50		
55		

	Ser Ala Gly Glu Lys Ala Pro Ser Pro Trp Gly Ser Ile Arg Thr	
5	830 835 840	
	ggg gct caa gct gca cac gtg tgg acc cct gcg gca ggg tcg tgc	2565
	Gly Ala Gln Ala Ala His Val Trp Thr Pro Ala Ala Gly Ser Cys	
10	845 850 855	
	tcc gtc tcc tgc ggg cga ggt ctg atg gag ctg cgt ttc ctg tgc	2610
	Ser Val Ser Cys Gly Arg Gly Leu Met Glu Leu Arg Phe Leu Cys	
15	860 865 870	
	atg gac tct gcc ctc agg gtg cct gtc cag gaa gag ctg tgt ggc	2655
20	Met Asp Ser Ala Leu Arg Val Pro Val Gln Glu Glu Leu Cys Gly	
	875 880 885	
25	ctg gca agc aag cct ggg agc cgg cgg gag gtc tgc cag gct gtc	2700
	Leu Ala Ser Lys Pro Gly Ser Arg Arg Glu Val Cys Gln Ala Val	
	890 895 900	
30	ccg tgc cct gct cgg tgg cag tac aag ctg gcg gcc tgc agc gtg	2745
	Pro Cys Pro Ala Arg Trp Gln Tyr Lys Leu Ala Ala Cys Ser Val	
	905 910 915	
35	agc tgt ggg aga ggg gtc gtg cgg agg atc ctg tat tgt gcc cgg	2790
	Ser Cys Gly Arg Gly Val Val Arg Arg Ile Leu Tyr Cys Ala Arg	
40	920 925 930	
	gcc cat ggg gag gac gat ggt gag gag atc ctg ttg gac acc cag	2835
	Ala His Gly Glu Asp Asp Gly Glu Glu Ile Leu Leu Asp Thr Gln	
45	935 940 945	
	tgc cag ggg ctg cct cgc ccg gaa ccc cag gag gcc tgc agc ctg	2880
	Cys Gln Gly Leu Pro Arg Pro Glu Pro Gln Glu Ala Cys Ser Leu	
50	950 955 960	
	gag ccc tgc cca cct agg tgg aaa gtc atg tcc ctt ggc cca tgt	2925
55	Glu Pro Cys Pro Pro Arg Trp Lys Val Met Ser Leu Gly Pro Cys	

	965	970	975	
5	tcg gcc agc tgt ggc ctt ggc act gct aga cgc tgc gtg gcc tgt			2970
	Ser Ala Ser Cys Gly Leu Gly Thr Ala Arg Arg Ser Val Ala Cys			
	980	985	990	
10	glg cag ctc gac caa ggc cag gac gtg gag gtg gac gag gcg gcc			3015
	Val Gln Leu Asp Gln Gly Gln Asp Val Glu Val Asp Glu Ala Ala			
	995	1000	1005	
15	tgt gcg gcg ctg glg cgg ccc gag gcc agt gtc ccc tgt ctc att			3060
	Cys Ala Ala Leu Val Arg Pro Glu Ala Ser Val Pro Cys Leu Ile			
	1010	1015	1020	
20	gcc gac tgc acc tac cgc tgg cat gtt ggc acc tgg atg gag tgc			3105
	Ala Asp Cys Thr Tyr Arg Trp His Val Gly Thr Trp Met Glu Cys			
	1025	1030	1035	
25	tct gtt tcc tgt ggg gat ggc atc cag cgc cgg cgt gac acc tgc			3150
	Ser Val Ser Cys Gly Asp Gly Ile Gln Arg Arg Arg Asp Thr Cys			
	1040	1045	1050	
30	ctc gga ccc cag gcc cag gcg cct gtg cca gct gat ttc tgc cag			3195
	Leu Gly Pro Gln Ala Gln Ala Pro Val Pro Ala Asp Phe Cys Gln			
	1055	1060	1065	
35	cac ttg ccc aag ccg gtg act gtg cgt ggc tgc tgg gct ggg ccc			3240
	His Leu Pro Lys Pro Val Thr Val Arg Gly Cys Trp Ala Gly Pro			
	1070	1075	1080	
40	tgt gtg gga cag ggt gcc tgt ggc agg cag cac ctt gag cca aca			3285
	Cys Val Gly Gln Gly Ala Cys Gly Arg Gln His Leu Glu Pro Thr			
	1085	1090	1095	
45	gga acc att gac atg cga ggc cca ggg cag gca gac tgt gca gtg			3330
	Gly Thr Ile Asp Met Arg Gly Pro Gly Gln Ala Asp Cys Ala Val			
	1100	1105	1110	
50				
55				

	gcc att ggg cgg ccc ctc ggg gag gtg gtg acc ctc cgc gtc ctt	3375
5	Ala Ile Gly Arg Pro Leu Gly Glu Val Val Thr Leu Arg Val Leu	
	1115 1120 1125	
	gag agt tct ctc aac tgc agt gcg ggg gac atg ttg ctg ctt tgg	3420
10	Glu Ser Ser Leu Asn Cys Ser Ala Gly Asp Met Leu Leu Leu Trp	
	1130 1135 1140	
15	ggc cgg ctc acc tgg agg aag atg tgc agg aag ctg ttg gac atg	3465
	Gly Arg Leu Thr Trp Arg Lys Met Cys Arg Lys Leu Leu Asp Met	
	1145 1150 1155	
20	act ttc agc tcc aag acc aac acg ctg gtg gtg agg cag cgc tgc	3510
	Thr Phe Ser Ser Lys Thr Asn Thr Leu Val Val Arg Gln Arg Cys	
	1160 1165 1170	
25	ggg cgg cca gga ggt ggg gtg ctg ctg cgg tat ggg agc cag ctt	3555
	Gly Arg Pro Gly Gly Gly Val Leu Leu Arg Tyr Gly Ser Gln Leu	
30	1175 1180 1185	
	gct cct gaa acc ttc tac aga gaa tgt gac atg cag ctc ttt ggg	3600
	Ala Pro Glu Thr Phe Tyr Arg Glu Cys Asp Met Gln Leu Phe Gly	
35	1190 1195 1200	
	ccc tgg ggt gaa atc gtg agc ccc tcg ctg agt cca gcc acg agt	3645
40	Pro Trp Gly Glu Ile Val Ser Pro Ser Leu Ser Pro Ala Thr Ser	
	1205 1210 1215	
	aat gca ggg ggc tgc cgg ctc ttc att aat gtg gct ccg cac gca	3690
45	Asn Ala Gly Gly Cys Arg Leu Phe Ile Asn Val Ala Pro His Ala	
	1220 1225 1230	
50	cgg att gcc atc cat gcc ctg gcc acc aac atg ggc gct ggg acc	3735
	Arg Ile Ala Ile His Ala Leu Ala Thr Asn Met Gly Ala Gly Thr	
	1235 1240 1245	
55	gag gga gcc aat gcc agc tac atc ttg atc cgg gac acc cac agc	3780

	Glu Gly Ala Asn Ala Ser Tyr Ile Leu Ile Arg Asp Thr His Ser	
5	1250 1255 1260	
	ttg agg acc aca gcg ttc cat ggg cag cag gtg ctg tac tgg gag	3825
10	1265 1270 1275	
	Leu Arg Thr Thr Ala Phe His Gly Gln Gln Val Leu Tyr Trp Glu	
15	1280 1285 1290	
	tca gag agc agc cag gct gag atg gag ttc agc gag ggc ttc ctg	3870
	Ser Glu Ser Ser Gln Ala Glu Met Glu Phe Ser Glu Gly Phe Leu	
20	1295 1300 1305	
	aag gct cag gcc agc ctg cgg ggc cag tac tgg acc ctg caa tca	3915
	Lys Ala Gln Ala Ser Leu Arg Gly Gln Tyr Trp Thr Leu Gln Ser	
25	1310 1315 1320	
	tgg gta ccg gag atg cag gac cct cag tcc tgg aag gga aag gaa	3960
	Trp Val Pro Glu Met Gln Asp Pro Gln Ser Trp Lys Gly Lys Glu	
30		3966
	gga acc	
	Gly Thr	
35	<210>20	
	<211>312	
40	<212>PRT	
	<213> Homo sapiens	
	<400>20	
45	gct gca ggc ggc atc cta cac ctg gag ctg ctg gtg gcc gtg ggc	45
	Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly	
50	1 5 10 15	
	ccc gat gtc ttc cag gct cac cag gag gac aca gag cgc tat gtg	90
	Pro Asp Val Phe Gln Ala His Gln Glu Asp Thr Glu Arg Tyr Val	
55	20 25 30	

	ctc acc aac ctc aac atc ggg gca gaa ctg ctt cgg gac ccg tcc	135
5	Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro Ser	
	35 40 45	
10	ctg ggg gct cag ttt cgg gtg cac ctg gtg aag atg gtc att ctg	180
	Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile Leu	
	50 55 60	
15	aca gag cct gag ggt gct cca aat atc aca gcc aac ctc acc tgc	225
	Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr Ser	
	65 70 75	
20	tcc ctg ctg agc gtc tgt ggg tgg agc cag acc atc aac cct gag	270
	Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro Glu	
	80 85 90	
25	gac gac acg gat cct ggc cat gct gac ctg gtc ctc tat atc act	315
	Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile Thr	
30	95 100 105	
	agg ttt gac ctg gag ttg cct gat ggt aac cgg cag gtg cgg ggc	360
	Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg Gly	
35	110 115 120	
	gtc acc cag ctg ggc ggt gcc tgc tcc cca acc tgg agc tgc ctc	405
40	Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys Leu	
	125 130 135	
	att acc gag gac act ggc ttc gac ctg gga gtc acc att gcc cat	450
45	Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala His	
	140 145 150	
50	gag att ggg cac agc ttc ggc ctg gag cac gac ggc gcg ccc ggc	495
	Glu Ile Gly His Ser Phe Gly Leu Glu His Asp Gly Ala Pro Gly	
	155 160 165	
55	agc ggc tgc ggc ccc agc gga cac gtg atg gct tgc gac ggc gcc	540

	Ser Gly Cys Gly Pro Ser Gly His Val Met Ala Ser Asp Gly Ala	
5	170 175 180	
	gcg ccc cgc gcc ggc ctc gcc tgg tcc ccc tgc agc cgc cgg cag	585
	Ala Pro Arg Ala Gly Leu Ala Trp Ser Pro Cys Ser Arg Arg Gln	
10	185 190 195	
	ctg ctg agc ctg ctc agg acg ggc gcg ctg cgt gtg gga ccc gcc	630
15	200 205 210	
	Leu Leu Ser Leu Leu Arg Thr Gly Ala Leu Arg Val Gly Pro Ala	
	gcg gcc tca acc cgg gtc cgc ggg gca ccc gcc gga tgc gca gcc	675
20	215 220 225	
	Ala Ala Ser Thr Arg Val Arg Gly Ala Pro Ala Gly Cys Ala Ala	
	tgg cct cta cta cag cgc caa cga gca gtg ccg cgt ggc ctt cgg	720
25	230 235 240	
	Trp Pro Leu Leu Gln Arg Gln Arg Ala Val Pro Arg Gly Leu Arg	
	ccc caa ggc tgt cgc ctg cac ctt cgc cag gga gca cct gga tat	765
30	245 250 255	
	Pro Gln Gly Cys Arg Leu His Leu Arg Gln Gly Ala Pro Gly Tyr	
	gtg cca ggc cct ctc ctg cca cac aga ccc gct gga cca aag cag	810
35	260 265 270	
	Val Pro Gly Pro Leu Leu Pro His Arg Pro Ala Gly Pro Lys Gln	
	ctg cag ccg cct cct cgt tcc tct cct gga tgg gac aga atg tgg	855
40	275 280 285	
	Leu Gln Pro Pro Pro Arg Ser Ser Pro Gly Trp Asp Arg Met Trp	
45	290 295 300	
	cgt gga gaa gtg gtg ctc caa ggg tcg ctg ccg ctc cct ggt gga	900
50		
	Arg Gly Glu Val Val Leu Gln Gly Ser Leu Pro Leu Pro Gly Gly	
	gct gac ccc cat agc agc agt gca tgg gcg ctg gtc	936
55	Ala Asp Pro His Ser Ser Ser Ala Trp Ala Leu Val	

305

310

5

<210>21

<211>270

10

<212>PRT

<213> Homo sapiens

15

<400>21

gct gca ggc ggc atc cta cac ctg gag ctg ctg gtg gcc gtg ggc 45

Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val Gly

20

1 5 10 15

ccc gat gtc ttc cag gct cac cag gag gac aca gag cgc tat gtg 90

Pro Asp Val Phe Gln Ala His Gln Glu Asp Thr Glu Arg Tyr Val

25

20 25 30

ctc acc aac ctc aac atc ggg gca gaa ctg ctt cgg gac ccg tcc 135

Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro Ser

30

35 40 45

ctg ggg gct cag ttt cgg gtg cac ctg gtg aag atg gtc att ctg 180

Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile Leu

35

50 55 60

aca gag cct gag ggt gct cca aat atc aca gcc aac ctc acc tcg 225

Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr Ser

40

65 70 75

tcc ctg ctg agc gtc tgt ggg tgg agc cag acc atc aac cct gag 270

Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro Glu

45

80 85 90

gac gac acg gat cct ggc cat gct gac ctg gtc ctc tat atc act 315

Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile Thr

50

55

95 100 105

	agg ttt gac ctg gag tlg cct gat ggt aac cgg cag glg cgg ggc	360
5	Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg Gly	
	110 115 120	
	gtc acc cag ctg ggc ggt gcc tgc tcc cca acc tgg agc tgc ctc	405
10	Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys Leu	
	125 130 135	
	att acc gag gac act ggc ttc gac ctg gga gtc acc att gcc cat	450
15	Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala His	
	140 145 150	
	gag att ggg cac agc ttc ggc ctg gag cac gac ggc gcg ccc ggc	495
20	Glu Ile Gly His Ser Phe Gly Leu Glu His Asp Gly Ala Pro Gly	
	155 160 165	
	agc ggc tgc ggc ccc agc gga cac glg atg gct tgc gac ggc gcc	540
25	Ser Gly Cys Gly Pro Ser Gly His Val Met Ala Ser Asp Gly Ala	
	170 175 180	
	gcg ccc cgc gcc ggc ctc gcc tgg tcc ccc tgc agc cgc cgg cag	585
30	Ala Pro Arg Ala Gly Leu Ala Trp Ser Pro Cys Ser Arg Arg Gln	
	185 190 195	
	ctg ctg agc ctg ctc aga ccc gtc cct ccg tgc ccg ctc cct ctg	630
35	Leu Leu Ser Leu Leu Arg Pro Val Pro Pro Ser Pro Leu Pro Leu	
	200 205 210	
	ctg gcc acc cac ctc tgc gcc ggc agg agc ctt agt ctt ggt ccc	675
40	Leu Ala Thr His Leu Cys Ala Gly Arg Ser Leu Ser Leu Gly Pro	
	215 220 225	
	agc caa gag ccg gct cct ggt ggg ggg cgc ggg ccg aga act cct	720
45	Ser Gln Glu Pro Ala Pro Gly Gly Gly Arg Gly Pro Arg Thr Pro	
	230 235 240	
	gtt ccc act cac aaa agg cca cgc ttc caa acg ctt cca tcc tgc	765
50		
55		

Val Pro Thr His Lys Arg Pro Arg Phe Gln Thr Leu Pro Ser Ser
 245 250 255
 5 tgc cca ctc ctc cgt ccc gcc tcc tcc cgg tgt aca ccc cgg gac 810
 Cys Pro Leu Leu Arg Pro Ala Ser Ser Arg Cys Thr Pro Arg Asp
 10 260 265 270
 <210>22
 <211>43
 15 <212>DNA
 <213> Homo sapiens
 20 <400>22
 ggactcgagc caccaatgca ccagcgtcac ccccgggcaa gat 43
 25 <210>23
 <211>45
 30 <212>DNA
 <213> Homo sapiens
 <400>23
 35 tccgtcgact cattatcagg ttccttcctt tcccttcag gactg 45
 40 <210>24
 <211>30
 <212>DNA
 45 <213> Homo sapiens
 <400>24
 50 gglttgcaat gtagacactg gtcaggttgg 30
 <210>25
 55 <211>30

<212>DNA

<213> Homo sapiens

<400>25

ccaacctgac cagtgtctac attgccaacc

30

<210>26

<211>30

<212>DNA

<213> Homo sapiens

<400>26

ctttccacct aggtgggcag ggctccaggc

30

<210>27

<211>30

<212>DNA

<213> Homo sapiens

<400>27

gcctggagcc ctgcccacct aggtggaaag

30

<210>28

<211>33

<212>DNA

<213> Homo sapiens

<400>28

tcgagaaaaa gtctacgggg gcctaggttt tta

33

<210>29

<211>33

<212>DNA

<213> Homo sapiens

<400>29

agcttaaaaa cctaggcccc cgtagacttt ttc

33

<210>30

<211>30

<212>DNA

<213> Homo sapiens

<400>30

tcggccaatgg ccgcaggcgg catcctacac

30

<210>31

<211>28

<212>DNA

<213> Homo sapiens

<400>31

ggcaagctta tcagcggggc gcggcgcc

28

<210>32

<211>564

<212>DNA

<213> Homo sapiens

<400>32

ccatggccgc aggcggcatc ctacacctgg agctgctggt ggccgtgggc cccgatgtct 60

tccaggctca ccaggaggac acagagcgt atgtgtcac caacctcaac atcggggcag 120

aactgttcg ggacccglcc ctgggggctc agtttcgggt gcacctggig aagatggtca 180

ttctgacaga gcctgagggt gctccaaata tcacagccaa cctcacctcg tccctgtga 240

gcgtctgtgg gtaggagccag accatcaacc ctgaggacga cacggatcct ggccaigctg 300
 5 acctggtcct ctatatact aggtttgacc tggagtigcc tgaatgtaac cggcaggatgc 360
 ggggcgtcac ccagctgggc ggtgcctgct cccaacctg gagctgcctc attaccgagg 420
 aactggctt cgacctggga gtcacattg ccatgagat tgggcacagc ttcggcctgg 480
 10 agcacgacgg cgcgcccggc agcggctgcg gcccagcgg acacgtgatg gcttcggacg 540
 gcgccgcgcc ccgctgataa gctt 564

15

<210>33

<211>184

20

<212>PRT

<213> Homo sapiens

25

<400>33

Met Ala Ala Gly Gly Ile Leu His Leu Glu Leu Leu Val Ala Val
 1 5 10 15
 30 Gly Pro Asp Val Phe Gln Ala His Gln Glu Asp Thr Glu Arg Tyr
 20 25 30
 Val Leu Thr Asn Leu Asn Ile Gly Ala Glu Leu Leu Arg Asp Pro
 35 35 40 45
 Ser Leu Gly Ala Gln Phe Arg Val His Leu Val Lys Met Val Ile
 40 50 55 60
 Leu Thr Glu Pro Glu Gly Ala Pro Asn Ile Thr Ala Asn Leu Thr
 65 70 75
 45 Ser Ser Leu Leu Ser Val Cys Gly Trp Ser Gln Thr Ile Asn Pro
 80 85 90
 50 Glu Asp Asp Thr Asp Pro Gly His Ala Asp Leu Val Leu Tyr Ile
 95 100 105
 Thr Arg Phe Asp Leu Glu Leu Pro Asp Gly Asn Arg Gln Val Arg
 55 110 115 120

Gly Val Thr Gln Leu Gly Gly Ala Cys Ser Pro Thr Trp Ser Cys
 5 125 130 135
 Leu Ile Thr Glu Asp Thr Gly Phe Asp Leu Gly Val Thr Ile Ala
 10 140 145 150
 His Glu Ile Gly His Ser Phe Gly Leu Glu His Asp Gly Ala Pro
 15 155 160 165
 Gly Ser Gly Cys Gly Pro Ser Gly His Val Met Ala Ser Asp Gly
 170 175 180
 Ala Ala Pro Arg
 20 185

 <210>34
 <211>2529
 <212>DNA
 30 <213> Mus. musculus
 <400>34

 35 atgagccagc ttgacctgtg gttagcgtgc cagccttggt atgctgtcag 50
 gtgcagagga atcctcactg gtgccatctt cattctgggc tgctgggggc 100
 tctctgactt ccagaagagt ctcttcaag atctggagcc caaggatgtg 150
 40 tcttcttact ttggccacca tgctgtcca ttcacaggcc atcctccctc 200
 tcacctccag agactgagac ggagaaggac ttggaggac attctgcacc 250
 tggaactcct ggtagctgtg ggccccgatg ttccccgggc tcatcaggag 300
 45 gacacagaac gctacgtgct cactaatctc aatacgggt cagaactgtt 350
 gagaaaccca tccctgggag tccagttcca ggtgcacctg gtgaagctaa 400
 50 tcacctctc tgactcagag agtactccga atatcacggc caacatcacc 450
 tcatccttga tgagcgtctg cgagtggagc cagacgatca acccccacga 500
 55 tgacagggat ccaagtcacg ctgacctgat tctctatctc accagcaacg 550
 tggctgggtgc cactgtcctt gtagttcatt ttctcttctc aaggtttgac 600

	ctggagtlgc ctgatggcaa ccagcaggtt cggggltgca cccagctggg	650
5	aggtgccctgc tcccttccct ggagttgcc tttcacctgag galactggct	700
	ttgacctggg ggtcaccatc gcccatgaga ttgggcacag cttcgggctg	750
10	gaccatgatg gtgctccagg tagtggcagc acctgcaagg ccagtggcca	800
	cgtgatggcg gctgatggcg caacacctac tggagggacc ctggagtgg	850
	ctgccctgcag ccaaaggcag ttgcagcacc tactcagcac agggcaaagt	900
15	cactgtctcc aggaccacc tgggctgcag tcaggactta cacggcacca	950
	gctgatggca cagcctggcc tctactacag tgcagatgat cagtgccgtg	1000
20	tggcttccgg ttctggggct gtgcctgca cttctccag ggagggtctg	1050
	aacacagcac tcagtggctc ttccaccttg atccgtccg cagaccctg	1100
	ccagaagctc tggatggctc ctgaagctct caaattctcc ttctccacca	1150
25	aatccgacat ctggctctct ggctgcatca ttctagacat ggccacttgc	1200
	tccttccctga acgacacaga agccatgcaa ctgcggaagg ccatccgcca	1250
	tcateccagc agcctgaagc ccatectgaa aaccatggag gagaagcaaa	1300
30	tccttggtac agatgtctac tatttgcttc tgccttcat gttgcatac	1350
	aacccctccg atcgactggc aatcaaggat gtgatgcaag tcaccttcat	1400
35	gagcaactcc ttcaaaagct cctctgttgc gctgaatatg cagcggcaga	1450
	aggtcccat ctctatcact gacgtgctgc ttgaaggcaa catggccaac	1500
	atcttaggtg atggcagctg gctgtgtgtc tcccttctga acgacagcag	1550
40	gcactgtgac tcagggattg gctcgcagag acttgggttt gatcttcagt	1600
	cagtctcttg gacagagcac cctctgaaag atgtcatgca gaatttctcc	1650
45	agtcgaccag aggtccagct cagagccatt aacaagtgt tgacaatgcc	1700
	agaggaccag ctagcactgg caaaggacc agaagctgag atcccaagga	1750
	gcagttgat catctcttc ctgatggata ccttgcggag ccatectaac	1800
50	tctgaaaggc ttgttaatgt ggcttacaac gtgcttgcca ttatttccag	1850
	ccaaggacag atctcagaag agctggaaga ggaggggttg tticagcttg	1900
55	ccaagagaa cctggagcac ttccaagagg acagggacat ctgctctct	1950
	atcctgagcc tgcctggct cctcctggta gatgtgtca ctgtggacaa	2000

	agagcccttg gagcagctct ctggcatggt caccleggtg ctggclactc	2050
5	atccggagga cgtggaaata gcagaggctg gctglgcggt gctctggctg	2100
	ctgtccttgt tgggctgcat aaaggagagt cagtltgagc aggtggtagt	2150
	gctgctccig agaagcatcc agctgtgccc tggcagagta ctgctggta	2200
10	acaatgcalt ccgtggcttg gccagcctcg caaaggigtc cggcccaccc	2250
	tcacagttag agccaaatga ctgggtatcc agccccagcc cccitttgtg	2300
15	gaatcagaga ctacactatg tgaacaagca aaagctgttc atgcccttgt	2350
	gggtgctgag gcaagagcac cctcattact gctgtgctaa tgacctaca	2400
	tcagagcaca tccaggcagt actaagtga ctaaattgggt ttgaaaagaa	2450
20	gcacagttgt gtggaatctt gtgtggaatg tggctgcagg cagcaggaga	2500
	agaatatagg aggagcccca gggatttga	2529

25 <210>35

<211>2514

30 <212>DNA

<213> Mus musculus

<400>35

35	aggaagctcc caagagtaaa cactgcctga tgtcccgccc agccagcaag	50
	tgaacattgc aactaacca gaatcccagt cactagggct cctgtccggc	100
40	catcaactgc cttttctaaa gatgagccag ctttgcctgt ggttgacgtg	150
	ccagccttgt tatgtgtica gtgtcagagg aatcctcact ggtgccatct	200
	tcattctggg ctgctggggg ctctctgact tccagaagag tcttcttcaa	250
45	gatctggagc ccaaggatgt gtcttcttac ttggccacc atgtgtctcc	300
	attcacaggc catcctccct ctacctcca gagactgaga cggagaagga	350
50	ctttggagga cattctgcac ctggaactcc tggtagctgt gggccccgat	400
	gtttcccggg ctcatcagga ggacacagaa cgctacgtgc tctaatact	450
	caatatcggg tcagaactgt tgagaaaccc atccctggga gtccagtcc	500
55	aggtagacct ggtgaagcta atcacctct ctgactcaga gactactccg	550

	aatatcacgg ccaacatcac ctcatccttg atgagcgtct gcgagtggag	600
5	ccagacgata aacccccacg atgacagga tccaagtcac gctgacctga	650
	ttctctatat caccagggtt gacctggagt tgcctgatgg caaccagcag	700
	gttcggggtg tcaccagct gggagggtgcc tgcctccctt cctggagttg	750
10	ccttatcact gaggatactg gctttgacct gggggtcacc atcgcccatg	800
	agattgggca cagcttcggg ctggaccaatg atgggtgtcc aggtagtggc	850
15	agcacctgca aggccagtgg ccacgtgatg gcggctgatg gcgcaacacc	900
	tactggaggg accctggagt ggtctgcctg cagccaaagg cagttgcagc	950
	acctactcag cacagggcag atgcactgct tccaggaccc acctgggctg	1000
20	cagtcaggac ttacacggca ccagctgatg gcacagcctg gcctctacta	1050
	cagtcagat gatcagtgcc gtgtggcttt cggttctggg gctgtgcct	1100
25	gcaccttctc cagggagggt ctggatgtat gccaggccct gtcctgccac	1150
	acagaccccc tggaccaaag cagctgcagc cgcctccttg ttccctcct	1200
	ggatgggaca ggatgtgggt tggagaagtg gtgtcccaag gctcgtgtc	1250
30	gtccctagc tgagctggct cctgtggctg cagtacatgg acactggctt	1300
	agctggggcc cccatagtcc ctgctccga tctgtggag gagggtgat	1350
35	taccaggagg cggtaggtga acaaccacag gcctgcatit gggggacgtg	1400
	catgtgtggg tgaagacctc caggctaaga tgtgcaacac gcaggcttgt	1450
	gagaagactc agctggagtt catgtccgag cagtgtgccc agacagacag	1500
40	acaaccactg caactttccc aaggcactgc ctctttctac cactgggatg	1550
	ctgctgtgca gtatagtcaa ggagataccc tgtgcagaca catgtgctgg	1600
	gctgttggag aaagcttcat tgtcagccgt ggggacaggt tcttagatgg	1650
45	gacccttgt gtgccaagt gtccccagga tgatgggacc ctaagcctct	1700
	gtttgttggg cagctgcagg acctttggct gtgatggcag gatggactcc	1750
50	cagaagggtt gggatgcgtg ccagggtgtg ggaggagaca acagcacctg	1800
	cagctcacgg aatggttctt tcacagctgg gagagccaga gaatatgtca	1850
	cgttccatgat tgttactccc aacatgacca acgcacacat tgtcaaccgc	1900
55	aggcctctct tcacacactt ggcggtgagg atccagggcc actacattgt	1950

	ggcaggaag actagcatct caccacaacac cacctaccct tcccttcctg	2000
5	aggactaccg tctggaatac agagtgaac tcactgagga ccagctgccc	2050
	cacttagagg agattcacat ccggggaccc gtccgggatg acattgagat	2100
	tcaggigtac agacgatatg gaggagaata tggggatctt acacaccag	2150
10	acatcacctt ttcctacttt caactgaagc agcaggcagc ctgggtatgg	2200
	accgctaagc gtggaccctg ctcatgagc tctggggcag ggctgcgctg	2250
15	ggtgacctac agctgccagg atcaagctca agacaagtgg gtaaagaacg	2300
	cccagtcca agggagccca cagccacctg catggcaaga gccttgtgtc	2350
	tctgccccct gctccccata ttgggtagct ggggacttca gcccatgtag	2400
20	cgctgtctgt ggccggggcc ttcgggagcg gtactgcgc tgtgtagaga	2450
	cccaagatgg ctcttaaaag acactgccac ctgcccggcg cagagcagta	2500
25	gcccagcagc cagc	2514

30 <210>36

<211>3512

<212>DNA

35 <213> Mus musculus.

<400>36

40	aggaagctcc caagagtaaa cactgcctga tctcccgccc agccagcaag	50
	tgaacattgc aactaacca gaatcccagt cactagggct cctgtccggc	100
	catcaactgc ctttctaaa gatgagccag ctttgcctgt ggttgacgtg	150
45	ccagccttgt tatgtgtca gtgtcagagg aatcctcact ggtgccatct	200
	tcatctggg ctgtggggg ctctctgact tccagaagag tcttcttcaa	250
	gatctggagc ccaaggatgt gtcttcttac ttggccacc atgtgtctcc	300
50	attacaggc catctccct ctacctcca gagactgaga cggagaagga	350
	ctttggagga catctgcac ctggaactcc tggtagctgt gggccccgat	400
55	gtttccggg ctcatcagga ggacacagaa cgctacgtgc tactaatct	450

	caatatcggg tcagaactgt tgagaaaccc atccctggga gtccagttcc	500
5	aggatgcacct ggigaagctt atcaccctct ctgactcaga gactactccg	550
	aataatcacgg ccaacatcac ctcatccttg atgagcgtct gcgagtggag	600
	ccagacgatt aacccccacg atgacaggga tccaagtcac gctgacctga	650
10	ttctctatat caccagggtt gacctggagt tgcctgatgg caaccagcag	700
	gttcgggggtg tcaccagct gggagggtgc tgctcccttt cctggagtig	750
15	ccttatcact gaggatctg gctttgacct gggggtcacc atgccccatg	800
	agattgggca cagcttcggg ctggaccaig atggtgctcc aggtagtggc	850
	agcaccigca aggccagtgg ccacgtgatg gcggctgacg gcgcaacacc	900
20	cactggaggg accctggagt ggctgcccig cagccaaagg cagtgcagc	950
	acctactcag cacagggcaa atgcactgct tccaggacct acctgggctg	1000
25	cagtcaggac ttacacggca ccagctgatg gcacagccig gcctctacta	1050
	cagtgcagat gatcagtgcc gtgtggcttt cggttctggg gctgtgcct	1100
	gcaccttctc cagggagggt ctggatgtat gccaggccct gtctgccac	1150
30	acagaccct tggaccaaag cagctgcagc cgcctccttg ttctctctct	1200
	ggatgggaca gaatgtgggt tggagaagt gtgtccaag gctcgtgtc	1250
	gtcccttagc tgagctggct cctgtggctg cagtacatgg acactggctt	1300
35	agctggggcc cccatagtcc ctgctccga tccgtggag gaggttgat	1350
	taccaggagg cgggtgtgca acaacccag gcctgcattt gggggacgtg	1400
40	catgtgtggg tgaagacct caggctaaga tgtcaaacac gcaggcttgt	1450
	gagaagactc agctggagtt catgtccgag cagtgtccc agacagacag	1500
	acaaccactg caactttccc aaggcactgc ctcttctac cactgggatg	1550
45	ctgctgtgca gtaagtcaa ggagataccc tgtcagaca catgtgttg	1600
	gctgttggag aaagcttcat tgcagccgt ggggacaggt tccatgatg	1650
50	gacccttgt gtgccaagt gtctcagga tgatgggacc ctaagcctct	1700
	gtttgttggg cagctgcagg acctttggct gtgatggcag galggactcc	1750
	cagaaggitt gggatgcgt ccaggtgtgt ggaggagaca acagcacctg	1800
55	cagctcacgg aatggttctt tcacagctgg gagagccaga gaatatgtca	1850

	cgttcctgat tgttactccc aacatgacca acgcacacat tgtcaaccgc	1900
5	aggectctct tcacacactt ggcggtgagg atccagggcc actacattgt	1950
	ggcaggggaag actagcatct caccacaacac cacttacctt tcccttctgg	2000
	aggactaccg tgtggaatac agagtgacac tcactgagga ccagctgccc	2050
10	cacttagagg agattcacat ccggggaccc gtccgggatg acattgagat	2100
	tcagggtgac agacgatacg gaggagaata tggggatctt acacaccag	2150
15	acatcacctt ttcttacttt caactgaagc agcaggcagc ctgggtatgg	2200
	accgctaagc gtggaccctg ctcatgagc tgtggggcag ggctgcgctg	2250
	ggtgacctac agctgccagg atcaagctca agacaagtgg gtaaagaacg	2300
20	cccagtgcca agggagccca cagccacctg catggcaaga gccttgtgtc	2350
	tctgccccct gctccccata ttgggtagct ggggacttca gcccatgtag	2400
25	cgtgtcttgt ggcgggggcc ttccggagcg gtcactgcgc tgtgtagaga	2450
	cccaagatgg ctctttaaag aacttgccac ctgcccggcg cagagcagta	2500
	gcccagcagc cagcagcaga agtggaaaac tgcaactccc agccctgtcc	2550
30	caccaggtgg gaggtgtcag accctggccc ttgcatgcca tctgacctg	2600
	aggcaggtct ggactcaagg aatgtgacat gtgtgtccag ggcgggtgac	2650
	ccggagaagc cagaaactgc agggccctgc cgcaccgacg agatgtcagc	2700
35	tatgttgag cctgtctcca ggagcctgtg ttctccaggc ttgggtcagg	2750
	tggacaacac catgtctctg ggcgaggagg ctccatcccc ggtgggcagt	2800
40	gacaagccag gggtcaggc tgagcatgtg tggacccctc tgggtgggct	2850
	gtgtctccatc tcttgtggga gaggtctgaa ggaactgtat ttctgtgca	2900
	tggattctgt cctcaaaatg cctgtccagg aagagctatg cggcttggct	2950
45	agtaagcccc caagccggcg ggaggtctgc agggctcgcc cctgtcctgc	3000
	tcgggtgggag actcaagctt tggcaccgtg cccggtgacc tgtggtggg	3050
50	ggcgagtgcc actgtctgtt cgttgtgtgc agctagaccg tggccaccgc	3100
	atatctgtac ctactccaa gtgtctgcca gtgcctaagc caggctcctt	3150
	cgaggactgc agccctgagc ctgttcctgc lagggcacta gtgtgggaag	3200
55	ccgccccac attcgccgtc acaagatggc gctgacatcc tgtgttctaa	3250

gttagtaaac aaataatcig cgcatgagcc aagggatatt acgactactt 3300
 5 gtactctggt tttcccgtag acgtcagctc ggccatgggc tgcagccaat 3350
 cagggagtag tgcgtcctag gcaattggtg ttctctttta aatagaaggg 3400
 gtttcgtttt ttctcttttc ttgttcttta cactctggcc ccaaaaagat 3450
 10 gtaagcaata aagctttgcc gtaggaaaaa aaaaaaaaaa ggatccggtg 3500
 cctctagatc ag 3512

15

<210>37

20

<211>22

<212>PRT

25

<213> Homo sapiens

<400>37

Phe. Ser. Pro Ala. Pro Gln Pro Arg Arg. Leu Leu Pro Gly Pro Gln.

30

1 5 10 15

Glu Asn Ser Val Gln Ser Ser

35

20

40

<210>38

<211>30

<212>DNA

45

<213> Homo sapiens

<400>38

50

atgtgcaaca ctcaggcctg cgagaagacc 30

55

<210>39

<211>30

<212>DNA

<213> Homo sapiens

<400>39

ccaacctgac cagtgtctac attgccaacc

30

<210>40

<211>21

<212>DNA

<213> Homo sapiens

<400>40

ctggagccct gccacctag g

21

<210>41

<211>62

<212>DNA

<213> Homo sapiens

<400>41

gccgtcgact cttatcactt atcgtcacg tccttgtagt ctgacgat gaactccagc
tg

60

62

<210>42

<211>62

<212>DNA

<213> Homo sapiens

<400>42

5 gccgtcgact cttatcactt atcgtcacg tccttgiagt ccaggttggg ggtaactgtc 60
ag 62

10

<210>43

15 <211>62

<212>DNA

<213> Homo sapiens

20 <400>43

gccgtcgact cttatcactt atcgtcacg tccttgiagt ccacgtgtgc agcttgagcc 60
25 cc 62

30 <210>44

<211>62

<212>DNA

35 <213> Homo sapiens

<400>44

40 gccgtcgact cttatcactt atcgtcacg tccttgiagt ccctagggtgg gcagggctcc 60
ag 62

45

<210>45

50 <211>62

<212>DNA

<213> Homo sapiens

55 <400>45

gccgtcgact cttatcactt atcgatcatg tcctttagt caccctgtcc cacacagggc 60

5 CC 62

10 <210>46

<211>60

15 <212>DNA

<213> Homo sapiens

<400>46

20 tccaagcttg tcgactctta tcacttatcg tcatcgctct tgtatcggt tccttccttt 60

25 <210>47

<211>27

30 <212>DNA

<213> Artificial Sequence

35 <220>

<223> Description of Artificial Sequence: Synthetic DNA

40 <400>47

45 gactacaagg acgatgacga taagtga 27

<210>48

50 <211>8

<212>RPT

55 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400>47

Asp Tyr Lys Asp Asp Asp Asp Lys

1

5

Claims

1. A protease that is capable of cleaving a bond between residues Tyr-842 and Met-843 of von Willebrand factor (hereinafter referred to as "vWF") and comprises a polypeptide chain having the amino acid sequence Leu-Leu-Val-Ala-Val as a partial sequence or an amino acid sequence with deletion, substitution, or addition of one or several amino acids in said amino acid sequence.
2. The protease according to claim 1, which comprises a polypeptide chain having the amino acid sequence Ala-Ala-Gly-Gly-Ile-Leu-His-Leu-Glu-Leu-Leu-Val-Ala-Val as the N-terminal partial sequence of a mature protein or an amino acid sequence with deletion, substitution, or addition of one or several amino acids in said amino acid sequence.
3. The protease according to claim 1 or 2, which comprises a polypeptide chain having an amino acid sequence with deletion, substitution, or addition of one or several amino acids in the amino acid sequence as shown in SEQ ID NO: 3 or 7 or a partial sequence of any of the aforementioned amino acid sequences as the N-terminal partial sequence of a mature protein or the aforementioned amino acid sequence.
4. The protease according to any one of claims 1 to 3, which comprises a polypeptide chain having an amino acid sequence with deletion, substitution, or addition of one or several amino acids in the amino acid sequence as shown in any of SEQ ID NOs: 16 to 21.
5. The protease according to any one of claims 1 to 4, which has molecular weight of 105 to 160 kDa or 160 to 250 kDa in SDS-PAGE under reducing or non-reducing conditions.
6. A gene fragment encoding a protease that is capable of cleaving a bond between residues Tyr-842 and Met-843 of vWF and comprises a polypeptide chain having the amino acid sequence Leu-Leu-Val-Ala-Val as a partial sequence or an amino acid sequence with deletion, substitution, or addition of one or several amino acids in said amino acid sequence.
7. A gene fragment encoding the protease according to any one of claims 2 to 5.
8. DNA encoding the protease according to any one of claims 1 to 5, which comprises a nucleotide sequence encoding a polypeptide capable of cleaving a bond between residues Tyr 842 and Met 843 of vWF comprising CTG CTG GTG GCC GTG or with deletion, substitution, or addition of one or several nucleotides therein.
9. The DNA encoding a protease according to claim 8, which comprises a nucleotide sequence comprising GCT GCA GGC GGC ATC CTA CAC CTG GAG CTG CTG GTG GCC GTG or with deletion, substitution, or addition of one or several nucleotides therein.
10. The DNA encoding a protease according to claim 8 or 9, which comprises a nucleotide sequence with deletion, substitution or addition of one or several nucleotides in the nucleotide sequence as shown in SEQ ID NO: 6 or a partial sequence of any of the nucleotide sequences, or the nucleotide sequence.

11. The DNA encoding a protease according to any one of claims 8 to 10, which comprises a nucleotide sequence with deletion, substitution or addition of one or several nucleotides in the nucleotide sequence as shown in SEQ ID NO: 15 or a partial sequence of any of the nucleotide sequences, or the nucleotide sequence.
- 5 12. A vector comprising the DNA encoding a protease according to claim 8 or 9, which comprises a nucleotide sequence with deletion, substitution or addition of one or several nucleotides in the nucleotide sequence as shown in SEQ ID NO: 6 or 15 or a partial sequence of any of the nucleotide sequences, or the nucleotide sequence.
- 10 13. The vector according to claim 12 comprising a polypeptide encoding domain and specialized in the expression of said polypeptide.
14. A cell transformed or transfected with the vector according to claim 12.
- 15 15. A host cell transformed or transfected with the expression vector according to claim 13.
16. A pharmaceutical composition comprising the protease according to any one of claims 1 to 5.
- 20 17. The pharmaceutical composition according to claim 16, which is applied to treating diseases caused by deterioration in activity of the protease according to any one of claims 1 to 5, which is involved with gene defects or liver diseases.
- 25 18. The pharmaceutical composition according to claim 16 or 17, which is applied to the inhibition of platelet aggregation caused by the formation of excess vWF high-molecular-weight multimers.
- 29 19. The pharmaceutical composition according to claim 18, wherein the disease is thrombotic thrombocytopenic purpura.
- 30 20. An antibody against the protease according to any one of claims 1 to 5.
- 35 21. The antibody according to claim 20 against the protease according to any one of claims 1 to 5, which is capable of inhibiting or neutralizing the protease activity.
- 36 22. The antibody according to claim 20 against the protease according to any one of claims 1 to 5, which can be used for affinity purification of the protease.
- 40 23. A process for purifying the protease according to any one of claims 1 to 5, which utilizes the antibody according to claim 22.
- 45 24. A pharmaceutical composition or diagnostic agent comprising an antibody against the protease according to any one of claims 1 to 5.
25. An antagonist, inhibitor, agonist, or activity regulator against the protease according to any one of claims 1 to 5.
- 50 26. A pharmaceutical composition or diagnostic agent comprising an antagonist, inhibitor, agonist, or activity regulator against the protease according to any one of claims 1 to 5.
- 55 27. A pharmaceutical composition or diagnostic agent comprising the DNA according to any one of claims 8 to 11 or antisense DNA thereof.
28. The pharmaceutical composition according to claim 27, which is used for gene therapy intended to cure diseases caused by deterioration in activity of the protease according to any one of claims 1 to 5, which is involved with gene defects or liver diseases.
29. A process for assaying vWF-cleaving activity, wherein a protease-substrate reaction is carried out using vWF and vWF-cleaving protease on a membrane filter, and a substrate sample is then recovered from the filter, followed by SDS-PAGE analysis without Western blotting.
30. A process for screening for a compound capable of cleaving vWF, wherein the vWF-cleaving activity of a test

compound is assayed by the process according to claim 29.

31. A process for preparing the protease according to any one of claims 1 to 5, wherein human plasma fraction I paste is used as a starting material.
32. A homologue of the protease according to any one of claims 1 to 5 derived from a different animal species or a homologous protein thereof.
33. A gene encoding the homologue of the protein according to claim 32 derived from a different animal species or a homologous protein thereof.
34. An animal having a modified gene encoding the homologue of the protein according to claim 32 derived from a different animal species or a homologous protein thereof.

FIG 1




FIG 2




FIG 3




FIG 4




FIG 5A




FIG 5B




FIG 5C




FIG 6A




FIG 6B




FIG 6C




FIG 7




FIG 8A




FIG 8B





FIG 9

gct	gca	ggc	ggc	atc	cta	cac	ctg	gag	ctg	ctg	gtg	gcc	gtg	ggc
Ala	Ala	Gly	Gly	Ile	Leu	His	Leu	Glu	Leu	Leu	Val	Ala	Val	Gly
1				5					10					15
ccc	gat	gtc	ttc	cag	gct	cac	cag	aag	gac	aca	gag	cgc	tat	gtg
Pro	Asp	Val	Phe	Gln	Ala	His	Gln	Lys	Asp	Thr	Glu	Arg	Tyr	Val
				20					25					30
ctc	acc	aac	ctc	aac	atc	ggg	gca	gaa	ctg	ctt	cgg	gac	ccg	tcc
Leu	Thr	Asn	Leu	Asn	Ile	Gly	Ala	Glu	Leu	Leu	Arg	Asp	Pro	Ser
				35					40					45
ctg	ggg	gct	cag	ttt	cgg	gtg	cac	ctg	gtg	aag	atg	gtc	att	ctg
Leu	Gly	Ala	Gln	Phe	Arg	Val	His	Leu	Val	Lys	Met	Val	Ile	Leu
				50					55					60
aca	gag	cct	gag	ggg	gct	cca	aat	atc	aca	gca	aac	ctc	acc	tcc
Thr	Glu	Pro	Glu	Gly	Ala	Pro	Asn	Ile	Thr	Ala	Asn	Leu	Thr	Ser
				65					70					75
tcc	ctg	ctg	agc	gtc	tgt	ggg	tgg	agc	cag	acc	atc	aac	cct	gag
Ser	Leu	Leu	Ser	Val	Cys	Gly	Trp	Ser	Gln	Thr	Ile	Asn	Pro	Glu
				80					85					90
gac	gac	acg	gat	cct	ggc	cat	gct	gac	ctg	gtc	ctc	tat	atc	act
Asp	Asp	Thr	Asp	Pro	Gly	His	Ala	Asp	Leu	Val	Leu	Tyr	Ile	Thr
				95					100					105
agg	ttt	gac	ctg	gag	ttg	cct	gat	ggg	aac	cgg	cag	gtg	cgg	ggc
Arg	Phe	Asp	Leu	Glu	Leu	Pro	Asp	Gly	Asn	Arg	Gln	Val	Arg	Gly
				110					115					120
gtc	acc	cag	ctg	ggc	ggg	gcc	tgc	tcc	cca	acc	tgg	agc	tgc	ctc
Val	Thr	Gln	Leu	Gly	Gly	Ala	Cys	Ser	Pro	Thr	Trp	Ser	Cys	Leu
				125					130					135
att	acc	gag	gac	act	ggc	ttc	gac	ctg	gga	gtc	acc	att	gcc	cat
Ile	Thr	Glu	Asp	Thr	Gly	Phe	Asp	Leu	Gly	Val	Thr	Ile	Ala	His
				140					145					150
gag	att	ggg	cac	agc	ttc	ggc	ctg	gag	cac	gac				
Glu	Ile	Gly	His	Ser	Phe	Gly	Leu	Glu	His	Asp				
				155					160					

FIG 10



brain
heart
skeletal muscle
colon (no mucosa)
thymus
spleen

kidney
liver
small intestine
placenta
lung
peripheral blood leukocyte

FIG 11

gctgcaggcg gcatactaca cctggagctg ctggtagccg tggcccccga tgtcttcag

Primer 1

gctcaccaga aggacacaga gcgctatgtg ctcaccaacc tcaacatcgg ggcagaactg

Primer 3

cttcgggacc cgtccctggg ggctcagitt cgggtgcacc tggtagaat ggtcattctg

acagagcctg agggtagctc aaatatcaca gcaaacctca cctcgtcctt gctgagcgtc

tgtgggtgga gccagaccat caaccctgag gacgacacgg atcctggcca tgctgacctg

Primer 4

gtcctctata tcactaggtt tgacctggag ttgcctgatg gtaaccggca ggtgcggggc

gtcaccacagc tgggcggtag ctgctccca acctggagct gcctcattac cgaggacact

ggcttcgacc tgggagtcac cattgcccat gagattgggc acagcttcgg cctggagcac

Primer 2

gac

Primer 1

Sense: gctgcaggcg gcatactaca cctggagctg

Antisense : cagctccagg ttaggatgc cgcctgcagc

Primer 2

Sense : accattgccc atgagattgg g

Antisense : cccaatctca tgggcaatgg t

Primer 3

Sense : gcgctatgtg ctcaccaacc tcaacatcgg

Antisense : ccgatgttga ggttggtag cacatagcgc

Primer 4

Sense : atcaaccctg aggacgacac

Antisense : gtgtcgtcct cagggttat

FIG 12




FIG 13




FIG 14




FIG 15




FIG 16




FIG 17




FIG 18




FIG 19




FIG 20




FIG 21





FIG 22



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/04141

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.⁷ C12N15/57, C12N9/50, C12P21/00, A01K67/027, C12N1/15,
C12N1/19, C12N1/21, C12N15/00, A61K38/46, A61P7/02, A61P43/00,
A61K45/00, A61K48/00, A61K31/711, G01N33/573.A, G01N33/573.Z,
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.⁷ C12N15/00-15/57, C12N9/50, A61K38/46

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
SwissProt/PIR/GeneSeq, GenBank/EMBL/DDBJ/GeneSeq, BIOSIS (DIALOG)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X/Y/A	JP 2000-508918 A (Immuno AG.), 18 July, 2000 (18.07.00), & WO 97/41206 A3	1-5, 20-27, 29-32/ 16-19/6-15, 28, 33, 34
Y/A	Miha FURLAN et al., Acquired Deficiency of von Willebrand Factor-Cleaving Protease in a Patient With Thrombotic Thrombocytopenic Purpura., Blood, 15 April, 1998 (15.04.98), Vol.91, No.8, pages 2839 to 2846	16-19/ 1-15, 20-34

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:
"A" document defining the general state of the art which is not
considered to be of particular relevance
"E" earlier document but published on or after the international filing
date
"L" document which may throw doubts on priority claim(s) or which is
cited to establish the publication date of another citation or other
special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other
means
"P" document published prior to the international filing date but later
than the priority date claimed

"T" later document published after the international filing date or
priority date and not in conflict with the application but cited to
understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be
considered novel or cannot be considered to involve an inventive
step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be
considered to involve an inventive step when the document is
combined with one or more other such documents, such
combination being obvious to a person skilled in the art
"Z" document member of the same patent family

Date of the actual completion of the international search
14 June, 2002 (14.06.02)

Date of mailing of the international search report
02 July, 2002 (02.07.02)

Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/04141

Continuation of A. CLASSIFICATION OF SUBJECT MATTER
(International Patent Classification (IPC))

Int.Cl⁷ G01N33/15.Z, G01N33/50.Z

(According to International Patent Classification (IPC) or to both
national classification and IPC)

Form PCT/ISA/210 (extra sheet) (July 1998)